

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup>:</b> <b>C07D 307/91, 333/76, 307/93, A61K 31/34, 31/38, C07D 405/04, 209/88, C07C 311/21</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/06561</b> <b>(43) International Publication Date:</b> 10 February 2000 (10.02.00)
<b>(21) International Application Number:</b> PCT/US99/12273 <b>(22) International Filing Date:</b> 2 June 1999 (02.06.99)  <b>(30) Priority Data:</b> 60/095,006 30 July 1998 (30.07.98) US  <b>(71) Applicant (for all designated States except US):</b> WARNER-LAMBERT COMPANY [US/US]; 201 Tabor Road, Morris Plain, NJ 07950 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> O'BRIEN, Patrick, Michael [US/US]; 5501 Green Road, Stockbridge, MI 49285 (US). PICARD, Joseph, Armand [US/US]; 44809 Saltz Road, Canton, MI 48187 (US). SLISKOVIC, Drago, Robert [GB/US]; 3679 Hedgerow Drive, Saline, MI 48176 (US).  <b>(74) Agents:</b> RYAN, M., Andrea; Warner-Lambert Company, 201 Tabor Road, Morris Plains, NJ 07950 (US) et al.		<b>(81) Designated States:</b> AE, AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> TRICYCLIC SULFONAMIDES AND THEIR DERIVATIVES AS INHIBITORS OF MATRIX METALLOPROTEINASES  <b>(57) Abstract</b>  Tricyclic sulfonamide compounds and derivatives are described as well as methods for the preparation and pharmaceutical compositions of same, which are useful as inhibitors of matrix metalloproteinases, particularly gelatinase A, collagenase-3, and stromelysin-1 and for the treatment of multiple sclerosis, atherosclerotic plaque rupture, aortic aneurysm, heart failure, left ventricular dilation, restenosis, periodontal disease, corneal ulceration, treatment of burns, decubital ulcers, wound healing, cancer, inflammation, pain, arthritis, osteoporosis, renal disease, or other autoimmune or inflammatory disorders dependent upon tissue invasion by leukocytes or other activated migrating cells, acute and chronic neurodegenerative disorders including stroke, head trauma, spinal cord injury, Alzheimer's disease, amyotrophic lateral sclerosis, cerebral amyloid angiopathy, AIDS, Parkinson's disease, Huntington's disease, prion diseases, myasthenia gravis, and Duchenne's muscular dystrophy.  <div style="text-align: right; margin-top: 200px;"><b>Document AW</b> <b>Cited in IDS for UCSF-085CIP</b> <b>Serial No. 09/616,223</b> <b>filed July 14, 2000</b></div>		



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						



-1-

## TRICYCLIC SULFONAMIDES AND THEIR DERIVATIVES AS INHIBITORS OF MATRIX METALLOPROTEINASES

### BACKGROUND OF THE INVENTION

The present invention relates to novel tricyclic sulfonamide compounds  
5 and their derivatives useful as pharmaceutical agents, to methods for their  
production, to pharmaceutical compositions which include these compounds and a  
pharmaceutically acceptable carrier, and to pharmaceutical methods of treatment.  
The novel compounds of the present invention are inhibitors of matrix  
metalloproteinases, e.g., gelatinase A (MMP-2), collagenase-3 (MMP-13), and  
10 stromelysin-1 (MMP-3). More particularly, the novel compounds of the present  
invention are useful in the treatment of atherosclerotic plaque rupture, aortic  
aneurism, heart failure, left ventricular dilation, restenosis, periodontal disease,  
corneal ulceration, treatment of burns, decubital ulcers, wound repair, cancer,  
inflammation, pain, arthritis, osteoporosis, multiple sclerosis, renal disease, and  
15 other autoimmune or inflammatory disorders dependent on the tissue invasion of  
leukocytes or other activated migrating cells. Additionally, the compounds of the  
present invention are useful in the treatment of acute and chronic  
neurodegenerative disorders including stroke, head trauma, spinal cord injury,  
Alzheimer's disease, amyotrophic lateral sclerosis, cerebral amyloid angiopathy,  
20 AIDS, Parkinson's disease, Huntington's disease, prion diseases, myasthenia  
gravis, and Duchenne's muscular dystrophy.

Gelatinase A and stromelysin-1 are members of the matrix  
metalloproteinase (MMP) family (Woessner J.F., *FASEB J.*, 1991;5:2145-2154).  
Other members include fibroblast collagenase, neutrophil collagenase,  
25 gelatinase B (92 kDa gelatinase), stromelysin-2, stromelysin-3, matrilysin,  
collagenase 3 (Freije J.M., Diez-Itza I., Balbin M., Sanchez L.M., Blasco R.,  
Tolivia J., and Lopez-Otin C., *J. Biol. Chem.*, 1994;269:16766-16773), and the  
newly discovered membrane-associated matrix metalloproteinases (Sato H.,  
Takino T., Okada Y., Cao J., Shinagawa A., Yamamoto E., and Seiki M., *Nature*,  
30 1994;370:61-65).



-2-

The catalytic zinc in matrix metalloproteinases is a focal point for inhibitor design. The modification of substrates by introducing chelating groups has generated potent inhibitors such as peptide hydroxymates and thiol-containing peptides. Peptide hydroxamates and the natural endogenous inhibitors of MMPs (TIMPs) have been used successfully to treat animal models of cancer and inflammation.

The ability of the matrix metalloproteinases to degrade various components of connective tissue makes them potential targets for controlling pathological processes. For example, the rupture of an atherosclerotic plaque is the most common event initiating coronary thrombosis. Destabilization and degradation of the extracellular matrix surrounding these plaques by MMPs has been proposed as a cause of plaque fissuring. The shoulders and regions of foam cell accumulation in human atherosclerotic plaques show locally increased expression of gelatinase B, stromelysin-1, and interstitial collagenase. In situ zymography of this tissue revealed increased gelatinolytic and caseinolytic activity (Galis Z.S., Sukhova G.K., Lark M.W., and Libby P., "Increased expression of matrix metalloproteinases and matrix degrading activity in vulnerable regions of human atherosclerotic plaques," *J. Clin. Invest.*, 1994;94:2494-2503). In addition, high levels of stromelysin RNA message have been found to be localized to individual cells in atherosclerotic plaques removed from heart transplant patients at the time of surgery (Henney A.M., Wakeley P.R., Davies M.J., Foster K., Hembry R., Murphy G., and Humphries S., "Localization of stromelysin gene expression in atherosclerotic plaques by in situ hybridization," *Proc. Nat'l. Acad. Sci.*, 1991;88:8154-8158).

Inhibitors of matrix metalloproteinases will have utility in treating degenerative aortic disease associated with thinning of the medial aortic wall. Increased levels of the proteolytic activities of MMPs have been identified in patients with aortic aneurisms and aortic stenosis (Vine N. and Powell J.T., "Metalloproteinases in degenerative aortic diseases," *Clin. Sci.*, 1991;81:233-239).

Heart failure arises from a number of diverse etiologies, but a common characteristic is cardiac dilation, which has been identified as an independent risk factor for mortality (Lee T.H., Hamilton M.A., Stevenson L.W., Moriguchi J.D., Fonarow G.C., Child J.S., Laks H., and Walden J.A., "Impact of left ventricular



size on the survival in advanced heart failure," *Am. J. Cardiol.*, 1993;72:672-676). This remodeling of the failing heart appears to involve the breakdown of extracellular matrix. Matrix metalloproteinases are increased in patients with both idiopathic and ischemic heart failure (Reddy H.K., Tyagi S.C., Tjaha I.E.,  
5 Voelker D.J., Campbell S.E., and Weber K.T., "Activated myocardial collagenase in idiopathic dilated cardiomyopathy," *Clin. Res.*, 1993;41:660A; Tyagi S.C., Reddy H.K., Voelker D., Tjaha I.E., and Weber K.T., "Myocardial collagenase in failing human heart," *Clin. Res.*, 1993;41:681A). Animal models of heart failure have shown that the induction of gelatinase is important in cardiac dilation  
10 (Armstrong P.W., Moe G.W., Howard R.J., Grima E.A., and Cruz T.F., "Structural remodeling in heart failure: gelatinase induction," *Can. J. Cardiol.*, 1994;10:214-220), and cardiac dilation precedes profound deficits in cardiac function (Sabbah H.N., Kono T., Stein P.D., Mancini G.B., and Goldstein S., "Left ventricular shape changes during the course of evolving heart failure," *Am.*  
15 *J. Physiol.*, 1992;263:H266-270).

Neointimal proliferation, leading to restenosis, frequently develops after coronary angioplasty. The migration of vascular smooth muscle cells (VSMCs) from the tunica media to the neointima is a key event in the development and progression of many vascular diseases and a highly predictable consequence of  
20 mechanical injury to the blood vessel (Bendeck M.P., Zempo N., Clowes A.W., Galaray R.E., and Reidy M., "Smooth muscle cell migration and matrix metalloproteinase expression after arterial injury in the rat," *Circulation Research*, 1994;75:539-545). Northern blotting and zymographic analyses indicated that gelatinase A was the principal MMP expressed and excreted by these cells.  
25 Further, antisera capable of selectively neutralizing gelatinase A activity also inhibited VSMC migration across basement membrane barrier. After injury to the vessel, gelatinase A activity increased more than 20-fold as VSMCs underwent the transition from a quiescent state to a proliferating, motile phenotype (Pauly R.R., Passaniti A., Bilato C., Monticone R., Cheng L., Papadopoulos N.,  
30 Gluzband Y.A., Smith L., Weinstein C., Lakatta E., and Crow M.T., "Migration of cultured vascular smooth muscle cells through a basement membrane barrier requires type IV collagenase activity and is inhibited by cellular differentiation," *Circulation Research*, 1994;75:41-54).



- Collagenase and stromelysin activities have been demonstrated in fibroblasts isolated from inflamed gingiva (Uitto V.J., Applegren R., and Robinson P.J., "Collagenase and neutral metalloproteinase activity in extracts from inflamed human gingiva," *J. Periodontal Res.*, 1981;16:417-424), and enzyme levels have been correlated to the severity of gum disease (Overall C.M., Wiebkin O.W., and Thonard J.C., "Demonstrations of tissue collagenase activity in vivo and its relationship to inflammation severity in human gingiva," *J. Periodontal Res.*, 1987;22:81-88). Proteolytic degradation of extracellular matrix has been observed in corneal ulceration following alkali burns (Brown S.I., Weller C.A., and Wasserman H.E., "Collagenolytic activity of alkali burned corneas," *Arch. Ophthalmol.*, 1969;81:370-373). Thiol-containing peptides inhibit the collagenase isolated from alkali-burned rabbit corneas (Burns F.R., Stack M.S., Gray R.D., and Paterson C.A., *Invest. Ophthalmol.*, 1989;30:1569-1575).
- Stromelysin is produced by basal keratinocytes in a variety of chronic ulcers (Saarialho-Kere U.K., Ulpu K., Pentland A.P., Birkedal-Hansen H., Parks W.O., and Welgus H.G., "Distinct Populations of Basal Keratinocytes Express Stromelysin-1 and Stromelysin-2 in Chronic Wounds," *J. Clin. Invest.*, 1994;94:79-88).
- Stromelysin-1 mRNA and protein were detected in basal keratinocytes adjacent to but distal from the wound edge in what probably represents the sites of the proliferating epidermis. Stromelysin-1 may thus prevent the epidermis from healing.
- Davies et al., (*Cancer Res.*, 1993;53:2087-2091) reported that a peptide hydroxymate, BB-94, decreased the tumor burden and prolonged the survival of mice bearing human ovarian carcinoma xenografts. A peptide of the conserved MMP propeptide sequence was a weak inhibitor of gelatinase A and inhibited human tumor cell invasion through a layer of reconstituted basement membrane (Melchiori A., Albili A., Ray J.M., and Stetler-Stevenson W.G., *Cancer Res.*, 1992;52:2353-2356). The natural tissue inhibitor of metalloproteinase-2 (TIMP-2) also showed blockage of tumor cell invasion in in vitro models (DeClerck Y.A., Perez N., Shimada H., Boone T.C., Langley K.E., and Taylor S.M., *Cancer Res.*, 1992;52:701-708). Studies of human cancers have shown that gelatinase A is



activated on the invasive tumor cell surface (Strongin A.Y., Marmer B.L., Grant G.A., and Goldberg G.I., *J. Biol. Chem.*, 1993;268:14033-14039) and is retained there through interaction with a receptor-like molecule (Monsky W.L., Kelly T., Lin C.-Y., Yeh Y., Stetler-Stevenson W.G., Mueller S.C., and  
5 Chen W.-T., *Cancer Res.*, 1993;53:3159-3164).

Inhibitors of MMPs have shown activity in models of tumor angiogenesis (Taraboletti G., Garofalo A., Belotti D., Drudis T., Borsotti P., Scanziani E., Brown P.D., and Giavazzi R., *Journal of the National Cancer Institute*, 1995;87:293 and Benelli R., Adatia R., Ensoli B., Stetler-Stevenson W.G.,  
10 Santi L., and Albini A, *Oncology Research*, 1994;6:251-257).

Several investigators have demonstrated consistent elevation of stromelysin and collagenase in synovial fluids from osteo- and rheumatoid arthritis patients as compared to controls (Walakovits L.A., Moore V.L., Bhardwaj N., Gallick G.S., and Lark M.W., "Detection of stromelysin and  
15 collagenase in synovial fluid from patients with rheumatoid arthritis and post-traumatic knee injury," *Arthritis Rheum.*, 1992;35:35-42; Zafarullah M., Pelletier J.P., Cloutier J.M., and Marcel-Pelletier J., "Elevated metalloproteinases and tissue inhibitor of metalloproteinase mRNA in human osteoarthritic synovia," *J. Rheumatol.*, 1993;20:693-697). TIMP-1 and TIMP-2 prevented the formation of  
20 collagen fragments, but not proteoglycan fragments in both the bovine nasal and pig articular cartilage models for arthritis, while a synthetic peptide hydroxamate could prevent the formation of both fragments (Andrews H.J., Plumpton T.A., Harper G.P., and Cawston T.E., *Agents Actions*, 1992;37:147-154; Ellis A.J., Curry V.A., Powell E.K., and Cawston T.E., *Biochem. Biophys. Res. Commun.*,  
25 1994;201:94-101).

Gijbels et al., (*J. Clin. Invest.*, 1994;94:2177-2182) recently described a peptide hydroxamate, GM6001, that suppressed the development or reversed the clinical expression of experimental autoimmune encephalomyelitis (EAE) in a  
30 dose dependent manner, suggesting the use of MMP inhibitors in the treatment of autoimmune inflammatory disorders such as multiple sclerosis.

A recent study by Madri has elucidated the role of gelatinase A in the extravasation of T-cells from the blood stream during inflammation (Ramanic A.M., and Madri J.A., "The Induction of 72-kDa Gelatinase in T Cells



upon Adhesion to Endothelial Cells is VCAM-1 Dependent," *J. Cell Biology*, 1994;125:1165-1178). This transmigration past the endothelial cell layer is coordinated with the induction of gelatinase A and is mediated by binding to the vascular cell adhesion molecule-1 (VCAM-1). Once the barrier is compromised, edema and inflammation are produced in the CNS. Also, leukocytic migration across the blood-brain barrier is known to be associated with the inflammatory response in EAE. Inhibition of the metalloproteinase gelatinase A would block the degradation of extracellular matrix by activated T-cells that is necessary for CNS penetration.

These studies provide the basis for the expectation that an effective inhibitor of gelatinase A and/or stromelysin-1 would have value in the treatment of diseases involving disruption of extracellular matrix resulting in inflammation due to lymphocytic infiltration, inappropriate migration of metastatic or activated cells, or loss of structural integrity necessary for organ function.

Neuroinflammatory mechanisms are implicated in a broad range of acute and chronic neurodegenerative disorders, including stroke, head trauma, multiple sclerosis, and Alzheimer's disease, to name a few (McGeer E.G. and McGeer P.L., "Neurodegeneration and the immune system". In: Calne D.B., ed. *Neurodegenerative Diseases*, W.B. Saunders, 1994:277-300). Other disorders that may involve neuroinflammatory mechanisms include amyotrophic lateral sclerosis (Leigh P.N., "Pathogenic mechanisms in amyotrophic lateral sclerosis and other motor neuron disorders". In: Calne D.B., ed., *Neurodegenerative Diseases*, W.B. Saunders, 1994:473-88), cerebral amyloid angiopathy (Mandybur T.I. and Balko G., "Cerebral amyloid angiopathy with granulomatous angiitis ameliorated by steroid-cytosan treatment," *Clin. Neuropharm.*, 1992;15:241-7), AIDS (Gendelman H.E. and Tardieu M., "Macrophages/microglia and the pathophysiology of CNS injuries in AIDS," *J. Leukocyte Biol.*, 1994;56:387-8), Parkinson's disease, Huntington's disease, prion diseases, and certain disorders involving the peripheral nervous system, such as myasthenia gravis and Duchenne's muscular dystrophy. Neuroinflammation, which occurs in response to brain injury or autoimmune disorders, has been shown to cause destruction of healthy tissue (Martin R., MacFarland H.F., and McFarlin D.E., "Immunological aspects of demyelinating diseases," *Annu Rev. Immunol.*, 1992;10:153-87;



-7-

Clark R.K., Lee E.V., Fish C.J., et al., "Development of tissue damage, inflammation and resolution following stroke: an immunohistochemical and quantitative planimetric study," *Brain Res. Bull.*, 1993;31:565-72; Giulian D. and Vaca K., "Inflammatory glia mediate delayed neuronal damage after ischemia in the central nervous system," *Stroke*, 1993;24(Suppl 12):184-90; Patterson P.H., "Cytokines in Alzheimer's disease and multiple sclerosis," *Cur. Opinion Neurobiol.*, 1995;5:642-6; McGeer P.L., Rogers J., and McGeer E.G., "Neuroimmune mechanisms in Alzheimer disease pathogenesis," *Alzheimer Dis. Assoc. Disorders*, 1994;8:149-58; Martin R. and McFarland H.F., "Immunological aspects of experimental allergic encephalomyelitis and multiple sclerosis," *Crit. Rev. Clin. Lab. Sci.*, 1995;32:121-82; Rogers J., Webster S., Lue L.F., et al., "Inflammation and Alzheimer's disease pathogenesis". In: *Neurobiology of Aging*, 1996;17:681-686; Rothwell N.J. and Relton J.K., "Involvement of cytokines in acute neurodegeneration in the CNS," *Neurosci. Biobehav. Rev.*, 1993;17:217-27). The pathological profiles and clinical courses of these disorders differ widely, but they all have in common the participation of immune/inflammatory elements in the disease process. In particular, many neurodegenerative disorders are characterized by large numbers of reactive microglia in postmortem brain samples, indicative of an active inflammatory process (McGeer E.G. and McGeer P.L., supra., 1994).

Increasing attention is being directed toward inflammatory mechanisms in Alzheimer's disease. Several lines of evidence support the involvement of neuroinflammation in Alzheimer's disease: 1) There is a significant increase in inflammatory markers in the Alzheimer brain, including acute phase reactants, cytokines, complement proteins, and MHC molecules (McGeer et al., supra., 1994; Rogers et al., supra.); 2) There is evidence that  $\beta$ -amyloid induces neurodegenerative changes primarily through interactions with inflammatory molecules, and that inflammation alone is sufficient to induce neurodegeneration (Rogers et al., supra); and 3) Growing epidemiological data indicate that anti-inflammatory therapy can delay the onset and slow the progression of Alzheimer's disease (McGeer P.L. and Rogers J., "Anti-inflammatory agents as a therapeutic approach to Alzheimer's disease,"



*Neurology*, 1992;42:447-9; Canadian Study of Health and Aging, "Risk factors for Alzheimer's disease in Canada," *Neurology*, 1994;44:2073-80; Lucca U., Tettamanti M., Forloni G., and Spagnoli A., "Nonsteroidal antiinflammatory drug use in Alzheimer's disease," *Biol. Psychiatry*, 1994;36:854-66; Hampel H. and Müller N., "Inflammatory and immunological mechanisms in Alzheimer's disease," *DN&P*, 1995;8:599-608; Breitner J.C.S., Gau B.A., Welsh K.A., et al., "Inverse association of anti-inflammatory treatments and Alzheimer's disease: Initial results of a co-twin control study," *Neurology*, 1994;44:227-32; Breitner J.C.S., Welsh K.A., Helms M.J., et al., "Delayed onset of Alzheimer's disease with nonsteroidal anti-inflammatory and histamine H2 blocking drugs," *Neurobiol. Aging*, 1995;16:523-30; Andersen K., Launer L.J., Ott A., Hoes A.W., Breteler M.M.B., and Hofman A., "Do nonsteroidal anti-inflammatory drugs decrease the risk for Alzheimer's disease? The Rotterdam Study," *Neurology*, 1995;45:1441-5; Rich J.B., Rasmusson D.X., Folstein M.F., et al., "Nonsteroidal anti-inflammatory drugs in Alzheimer's disease," *Neurology*, 1995;45:51-5; Aisen P.S., "Anti-inflammatory therapy for Alzheimer's disease," *Dementia*, 1995;9:173-82; Rogers et al., supra). Chronic use of nonsteroidal anti-inflammatory drugs (NSAIDs), most commonly for the treatment of rheumatoid arthritis, decreases the probability of developing Alzheimer's disease, and there is reason to believe that other anti-inflammatory agents may also be effective, although direct evidence for the efficacy of such treatments is lacking (Hamper and Müller, supra., 1995). Furthermore, virtually all of the currently available compounds, which include corticosteroids, NSAIDs, antimalarial drugs, and colchicine, have serious drawbacks that make them undesirable in the treatment of chronic disorders. Glucocorticoids, which are in wide clinical use as anti-inflammatory/immunosuppressive drugs, can be directly neurotoxic and also are toxic to systemic organs at moderate to high doses. NSAIDs have gastrointestinal and renal side effects that obviate long-term use in most people, and few of them cross the blood-brain barrier in significant amounts. The toxic properties of chloroquine compounds and colchicine also are well known. An anti-inflammatory drug that is well-tolerated by patients and that crosses the blood-brain barrier has significant advantages for the treatment of acute and chronic degenerative diseases of the central nervous system.



Normal kidney function is dependent on the maintenance of tissues constructed from differentiated and highly specialized renal cells which are in a dynamic balance with their surrounding extracellular matrix (ECM) components (Davies M. et al., "Proteinases and glomerular matrix turnover," *Kidney Int.*, 1992;41:671-678). Effective glomerular filtration requires that a semi-permeable glomerular basement membrane (GBM) composed of collagens, fibronectin, enactin, laminin and proteoglycans is maintained. A structural equilibrium is achieved by balancing the continued deposition of ECM proteins with their degradation by specific metalloproteinases (MMP). The MMP belong to a supergene family of zinc endopeptidases (Woessner J.F., "Matrix metalloproteinases and their inhibitors in connective tissue remodelling," *FASEB J.*, 1991;5:2145-2154). These proteins are first secreted as proenzymes and are subsequently activated in the extracellular space. These proteinases are in turn subject to counter balancing regulation of their activity by naturally occurring inhibitors referred to as TIMPs (tissue inhibitors of metalloproteinases).

Deficiency or defects in any component of the filtration barrier may have catastrophic consequences for longer term renal function. For example, in hereditary nephritis of Alport's type, associated with mutations in genes encoding ECM proteins, defects in collagen assembly lead to progressive renal failure associated with splitting of the GBM and eventual glomerular and interstitial fibrosis. By contrast in inflammatory renal diseases such as glomerulonephritis, cellular proliferation of components of the glomerulus often precede obvious ultrastructural alteration of the ECM matrix. Cytokines and growth factors implicated in proliferative glomerulonephritis such as interleukin-1, tumor necrosis factor, and transforming growth factor beta can upregulate metalloproteinase expression in renal mesangial cells (Martin J. et al., "Enhancement of glomerular mesangial cell neutral proteinase secretion by macrophages: role of interleukin 1," *J. Immunol.*, 1986;137:525-529; Marti H.P. et al., "Homology cloning of rat 72 kDa type IV collagenase: Cytokine and second-messenger inducibility in mesangial cells," *Biochem. J.*, 1993;291:441-446; Marti H.P. et al., "Transforming growth factor-b stimulates glomerular mesangial cell synthesis of the 72 kDa type IV collagenase," *Am. J. Pathol.*, 1994;144:82-94). These metalloproteinases are believed to be intimately



involved in the aberrant tissue remodeling and cell proliferation characteristic of renal diseases, such as, IgA nephropathy which can progress to through a process of gradual glomerular fibrosis and loss of functional GBM to end-stage renal disease. Metalloproteinase expression has already been well-characterized in experimental immune complex-mediated glomerulonephritis such as the anti-Thy 1.1 rat model (Bagchus W.M., Hoedemaeker P.J., Rozing J., Bakker W.W., "Glomerulonephritis induced by monoclonal anti-Thy 1.1 antibodies: A sequential histological and ultrastructural study in the rat," *Lab. Invest.*, 1986;55:680-687; Lovett D.H., Johnson R.J., Marti H.P., Martin J., Davies M., Couser W.G., "Structural characterization of the mesangial cell type IV collagenase and enhanced expression in a model of immune complex mediated glomerulonephritis," *Am. J. Pathol.*, 1992;141:85-98).

Unfortunately, at present, there are very limited therapeutic strategies for modifying the course of progressive renal disease. Although many renal diseases have an inflammatory component, their responses to standard immunosuppressive regimes are unpredictable and potentially hazardous to individual patients. The secondary consequences of gradual nephron failure such as activation of the renin-angiotensin system, accompanied by individual nephron glomerular hyperfiltration and renal hypertension, may be effectively treated with ACE inhibitors or angiotensin II receptor antagonists; but at best, these compounds can only reduce the rate of GFR decline.

A novel strategy to treat at least some renal diseases has been suggested by recent observations of MMP behavior. A rat mesangial cell MMP has been cloned (MMP-2) which is regulated in a tissue specific manner, and in contrast to other cellular sources such as tumor cell lines, is induced by cytokines (Brown P.D., Levy A.T., Margulies I., Liotta L.A., Stetler-Stevenson W.G., "Independent expression and cellular processing of Mr 72,000 type IV collagenase and interstitial collagenase in human tumorigenic cell lines," *Cancer Res.*, 1990;50:6184-6191; Marti H.P. et al., "Homology cloning of rat 72 kDa type IV collagenase: Cytokine and second-messenger inducibility in mesangial cells," *Biochem. J.*, 1993;291:441-446). While MMP-2 can specifically degrade surrounding ECM, it also affects the phenotype of adjacent mesangial cells. Inhibition of MMP-2 by antisense oligonucleotides or transfection techniques can



-11-

induce a reversion of the proliferative phenotype of cultured mesangial cells to a quiescent or non-proliferative phenotype mimicking the natural in vitro behavior of these cells (Kitamura M. et al., "Gene transfer of metalloproteinase transin induces aberrant behaviour of cultured mesangial cells," *Kidney Int.*, 1994;45:1580-1586; Turck J. et al., "Matrix metalloproteinase 2 (gelatinase A) regulates glomerular mesangial cell proliferation and differentiation," *J. Biol. Chem.*, 1996;271:15074-15083).

Inhibitors of MMP (MMPi) clearly have potential clinical applications in a host of diseases characterized by disturbance of extracellular matrix-cell interactions resulting in abnormal tissue remodeling (Vincenti M.P. et al., "Using inhibitors of metalloproteinases to treat arthritis," *Arthritis Rheum.*, 1994;8:1115-1126; Grams F. et al., "X-ray structures of human neutrophil collagenase complexed with peptide hydroxamate and peptide thiol inhibitors. Implications for substrate binding and rational drug design," *Eur. J. Biochem.*, 1995;228:830-841).


We have identified a series of tricyclic sulfonamide compounds and their derivatives that are inhibitors of matrix metalloproteinases, particularly collagenase-3, stromelysin-1 and gelatinase A, and thus useful as agents for the treatment of multiple sclerosis, atherosclerotic plaque rupture, restenosis, aortic aneurism, heart failure, left ventricular dilation, periodontal disease, corneal ulceration, treatment of burns, decubital ulcers, wound repair, cancer, inflammation, pain, arthritis, osteoporosis, renal disease, or other autoimmune or inflammatory diseases dependent upon tissue invasion by leukocytes or other activated migrating cells, acute and chronic neurodegenerative disorders including stroke, head trauma, spinal cord injury, Alzheimer's disease, amyotrophic lateral sclerosis, cerebral amyloid angiopathy, AIDS, Parkinson's disease, Huntington's diseases, prion diseases, myasthenic gravis, and Duchenne's muscular dystrophy.

## SUMMARY OF THE INVENTION

Accordingly, a first aspect of the present invention is a compound of Formula I



-12-



wherein n is zero or an integer of 1 or 2;

X is -O-,

-S(O)<sub>p</sub>- wherein p is zero or an integer of 1 or 2,

5 -N- wherein R<sup>2</sup> is hydrogen,

|  
R<sup>2</sup>                      alkyl,  
                             acyl, or  
                             benzyl,

-CH<sub>2</sub>-, or

10 -C-;  
    ||  
    O

R is hydrogen,

alkyl,

15 hydroxyalkyl,

alkoxyalkyl,

trifluoromethyl,

alkanoyloxyalkyl,

alkanoylaminoalkyl,

20 alkylthioalkyl,

alkylsulfinylalkyl,

alkylsulfonylalkyl,

aminoalkyl,

alkylaminoalkyl,

25 dialkylaminoalkyl,

N-alkylpiperazinoalkyl,

N-phenylalkylpiperazinoalkyl,

morpholinoalkyl,

thiomorpholinoalkyl,

30 piperidinoalkyl,



-13-

- pyrrolidinoalkyl,  
N-alkylalkylpiperidinolalkyl,  
pyridylalkyl,  
thienylalkyl,  
5 quinolinyllalkyl,  
thiazolylalkyl,  
cycloalkyl,  
cycloalkylalkyl,  
phenyl,  
10 phenyl substituted by one to three substituents selected from the group  
consisting of:  
hydroxy,  
alkoxy,  
alkyl,  
15 alkylthio,  
alkylsulfinyl,  
alkylsulfonyl,  
amino,  
alkylamino,  
20 dialkylamino,  
halogen,  
cyano,  
nitro,  
trifluoromethyl or on adjacent carbon atoms by either a one  
25 to two carbon alkenylenedioxy group or a two to  
three carbon alkenyleneoxy group,  
phenylalkyl,  
phenylalkyl wherein phenyl is substituted by alkyl,  
alkoxy,  
30 halogen, or  
trifluoromethyl,  
heteroaryl,



-14-

heteroaryl substituted by one to two substituents selected from the group  
consisting of:

alkyl, or

halogen,

5 biphenyl,

biphenyl substituted by alkyl,

alkoxy,

halogen,

trifluoromethyl, or

10 cyano,

biphenylalkyl or

biphenylalkyl wherein biphenyl is substituted by

alkyl,

alkoxy,

15 halogen,

trifluoromethyl, or

cyano;

D is zero or an integer of 1 to 3;

L is zero or an integer of 1 to 3;

20  $R^1$  is hydrogen,

a side chain of a natural amino acid or

a side chain of an unnatural amino acid;

Y is  $OR^3$  wherein  $R^3$  is hydrogen,

methyl,

25 ethyl, or

benzyl, or

$NH-OR^4$  wherein  $R^4$  is hydrogen,

alkyl, or

benzyl;

30 and corresponding isomers thereof; or a pharmaceutically acceptable salt thereof.

As matrix metalloproteinase inhibitors, the compounds of Formula I are  
useful as agents for the treatment of multiple sclerosis. They are also useful as



agents for the treatment of atherosclerotic plaque rupture, aortic aneurism, heart failure, left ventricular dilation, restenosis, periodontal disease, corneal ulceration, treatment of burns, decubital ulcers, wound repair, cancer metastasis, tumor angiogenesis, inflammation, pain, arthritis, osteoporosis, renal disease, and other  
5 autoimmune or inflammatory disorders dependent upon tissue invasion by leukocytes or other activated migrating cells, acute and chronic neurodegenerative disorders including stroke, head trauma, spinal cord injury, Alzheimer's disease, amyotrophic lateral sclerosis, cerebral amyloid angiopathy, AIDS, Parkinson's disease, Huntington's disease, prion diseases, myasthenia gravis, and Duchenne's  
10 muscular dystrophy.

A still further embodiment of the present invention is a pharmaceutical composition for administering an effective amount of a compound of Formula I in unit dosage form in the treatment methods mentioned above. Finally, the present invention is directed to methods for production of compounds of Formula I.

#### 15 DETAILED DESCRIPTION OF THE INVENTION

In the compounds of Formula I, the term "alkyl" means a straight or branched hydrocarbon radical having from 1 to 6 carbon atoms and includes, for example, methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, sec-butyl, isobutyl, tert-butyl, *n*-pentyl, *n*-hexyl, and the like.

20 The term "alkenyl" means a straight or branched unsaturated hydrocarbon radical having from 2 to 6 carbon atoms and includes, for example, ethenyl, 2-propenyl, 1-butenyl, 2-butenyl, 1-pentenyl, 2-pentenyl, 3-methyl-3-butenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, and the like.


"Alkoxy" and "thioalkoxy" are O-alkyl or S-alkyl of from 1 to 6 carbon  
25 atoms as defined above for "alkyl".

The term "cycloalkyl" means a saturated hydrocarbon ring having 3 to 7 carbon atoms optionally containing an oxygen or sulfur atom and includes, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and the like.




-16-

The term "aryl" means an aromatic radical which is a phenyl group, a phenyl group substituted by 1 to 4 substituents selected from alkyl as defined above, alkoxy as defined above, thioalkoxy as defined above, hydroxy, halogen, trifluoromethyl, amino, alkylamino as defined above for alkyl, dialkylamino as defined above for alkyl, nitro, cyano, carboxy, guanidino, amidino, SO<sub>3</sub>H, CHO,



as defined above for alkyl,  $\begin{array}{c} \text{O} \\ || \\ \text{-C-N(alkyl)}_2 \end{array}$  as defined above for alkyl,  $\text{-(CH}_2\text{)}_{n^2}\text{-NH}_2$  wherein  $n^2$  is an integer of 1 to 5,  $\text{-(CH}_2\text{)}_{n^2}\text{-NH-alkyl}$  as defined above for alkyl and  $n^2$ ,  $\text{-(CH}_2\text{)}_{n^2}\text{-N(alkyl)}_2$  as defined above for alkyl and  $n^2$ ,



as defined above for alkyl and  $n^2$ .

The term "arylalkyl" means an aromatic radical attached to an alkyl radical wherein aryl and alkyl are as defined above for example benzyl, phenylethyl, 3-phenylpropyl, (4-chlorophenyl)methyl, and the like.

The term "acyloxymethyl" means a group of the formula  $\begin{array}{c} \text{O} \\ || \\ \text{-CH}_2\text{-O-C-alkyl} \end{array}$  wherein alkyl is as defined above.

The term "heteroaryl" means a 5- and 6-membered heteroaromatic radical containing 1 to 3 heteroatoms selected from N, O, and S and includes, for example, a heteroaromatic radical which is 2- or 3-thienyl, 2- or 3-furanyl, 2- or 3-pyrrolyl, 2-, 3-, or 4-pyridinyl, 2-pyrazinyl, 2-, 4-, or 5-pyrimidinyl, 3- or 4-pyridazinyl, 1H-indol-6-yl, 1H-indol-5-yl, 1H-benzimidazol-6-yl, 1H-benzimidazol-5-yl, 2-, 4-, or 5-thiazolyl, 3-, 4-, or 5-isothiazolyl, 2-, 4-, or 5-imidazolyl, 3-, 4-, or 5-pyrazolyl, or 2- or 5-thiadiazolyl optionally substituted by a substituent selected from alkyl as defined above, alkoxy as defined above,



-17-

thioalkoxy as defined above, hydroxy, halogen, trifluoromethyl, amino, alkylamino as defined above for alkyl, dialkylamino as defined above for alkyl, nitro, cyano, carboxy, guanidino, amidino,  $\text{SO}_3\text{H}$ ,  $\text{CHO}$ ,

5  $\begin{array}{c} \text{O} \\ \parallel \\ \text{C-alkyl} \end{array}$  as defined above for alkyl,  $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C-NH}_2 \end{array}$ ,  $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C-NH-alkyl} \end{array}$ ,  $\begin{array}{c} \text{O} \\ \parallel \\ \text{NH-C-alkyl} \end{array}$ ,

as defined above for alkyl,  $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C-N(alkyl)}_2 \end{array}$  as defined above for alkyl,  
10  $-(\text{CH}_2)_{n^2}\text{-NH}_2$  wherein  $n^2$  is an integer of 1 to 5,  $-(\text{CH}_2)_{n^2}\text{-NH-alkyl}$  as defined above for alkyl and  $n^2$ ,  $-(\text{CH}_2)_{n^2}\text{-N(alkyl)}_2$  as defined above for alkyl and  $n^2$ ,

15  $-(\text{CH}_2)_{n^2}\text{-NH-}\begin{array}{c} \text{O} \\ \parallel \\ \text{C-alkyl} \end{array}$  as defined above for alkyl, and  $n^2$  and  $-(\text{CH}_2)_{n^2}\text{-N-}\begin{array}{c} \text{O} \\ \parallel \\ \text{C-alkyl} \\ | \\ \text{alkyl} \end{array}$

as defined above for alkyl and  $n^2$ .

The term "heterocycle" means a 3- to 7-membered cycloalkyl radical containing 1 to 3 heteroatoms selected from N, O, and S and includes, for  
20 example, 2- and 3-azetidiny, 3- and 4-azetidiny-2-one, 4- and 5-imidazolidiny-2-one, 2,4-dioxo-imidazolidiny, 2,4-dioxo-1,5,5-trimethyl-imidazolidiny, 2-, 4-, and 5-thiazolidiny, 4- and 5-oxazolidiny-2-one, 2- and 3-tetrahydrofurany, 2- and 3-pyrrolidiny, 2-, 3-, and 4-piperidiny, 2- and 3-morpholiny, 2- and 3-piperaziny, 2-, 3-, and 4-azacycloheptany and the like.

25 The term "heteroarylalkyl" means a heteroaromatic radical attached to an alkyl radical wherein heteroaryl and alkyl are as defined above.

The term "heterocycloalkyl" means a heterocycle radical attached to an alkyl radical wherein heterocycle and alkyl are as defined above.

30 The term "pyrrolidinoalkyl" means a pyrrolidino group attached to an alkyl radical wherein alkyl is as defined above.

The term "pyridylalkyl" means a pyridyl group attached to an alkyl radical wherein alkyl is as defined above.

The term "thienylalkyl" means a thienyl group attached to an alkyl radical wherein alkyl is as defined above.



-18-

The term "quinolinylalkyl" means a quinolinyl group attached to an alkyl radical wherein alkyl is as defined above.

The term "thiazolylalkyl" means a thiazolyl group attached to an alkyl radical wherein alkyl is as defined above.

5 The term "phenylalkyl" means a phenyl group attached to an alkyl radical wherein alkyl is as defined above.

The term "biphenylalkyl" means a biphenyl group attached to an alkyl radical wherein alkyl is as defined above.

10 The term "acyl" means a group of the formula  $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}-\text{alkyl} \end{array}$  wherein alkyl is as defined above.

The term "hydroxyalkyl" means a hydroxy group attached to an alkyl radical wherein alkyl is as defined above.

15 The term "alkoxyalkyl" means an alkoxy group attached to an alkyl radical wherein alkoxy and alkyl are as defined above.

The term "aminoalkyl" means an amino group attached to an alkyl radical wherein alkyl is as defined above.

20 The term "morpholinoalkyl" means a morpholino group attached to an alkyl radical wherein alkyl is as defined above.

The term "thiomorpholinoalkyl" means a thiomorpholino group attached to an alkyl radical wherein alkyl is as defined above.

The term "piperidinoalkyl" means a piperidino group attached to an alkyl radical wherein alkyl is as defined above.

25 The term "cycloalkylalkyl" means a cycloalkyl group attached to an alkyl radical wherein cycloalkyl and alkyl are as defined above.

The terms "alkylaminoalkyl" and "dialkylaminoalkyl" are respectively alkyl-NH and (alkyl)<sub>2</sub>N- wherein alkyl is as defined above.

30 The term "alkylamino" and "dialkylamino" are respectively alkyl NH- and (alkyl)<sub>2</sub>N- wherein alkyl is as defined above.

The terms "alkylthioalkyl," "alkylsulfinylalkyl," and "alkylsulfonylalkyl" are respectively alkyl-S-alkyl, alkyl-SO-alkyl, and alkyl-SO<sub>2</sub>-alkyl wherein alkyl is as defined above.




The terms "alkylthio," "alkylsulfinyl," and "alkylsulfonyl" are respectively alkyl-S-, alkyl-SO-, and alkyl-SO<sub>2</sub>- wherein alkyl is as defined above.


5                   The term "alkanoyloxyalkyl" means an alkyl-C(=O)-O-alkyl wherein alkyl is as defined above.

10                   The term "alkanoylaminoalkyl" means an alkyl-C(=O)-NH-alkyl wherein alkyl is as defined above.


The term "N-alkylpiperazinoalkyl" means



The term "N-phenylalkylpiperazinoalkyl" means



15                   The term "N-alkylalkylpiperidinoalkyl" means



The term "alkenylendioxy" means -O-alkyl-O- wherein alkyl is as defined above.

20                   The term "alkenyleneoxy" means -alkyl-O- wherein alkyl is as defined above.

The term "side chain of a natural amino acid" (natural α amino acid) means the group Q in a natural amino acid of formula H<sub>2</sub>N-CH(Q)-COOH.

Examples of side chains of natural α amino acids include those of alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glycine, histidine, 25 isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine.

A natural α amino acid is an amino acid found in a living organism. Examples of such amino acids include glycine, alanine, valine, leucine, isoleucine, phenylalanine, proline, serine, threonine, tyrosine, asparagine, glutamine, lysine,




-20-

arginine, tryptophan, histidine, cysteine, methionine, aspartic acid, and glutamic acid.

The functional groups in the amino acid side chains can be protected. For example, carboxyl groups can be esterified, amino groups can be converted to amides or carbamates, hydroxyl groups can be converted to ethers or esters, and thiol groups can be converted to thioethers or thioesters.

The term "side chain of an unnatural amino acid" means the group R<sup>1a</sup> in an unnaturally occurring amino acid of formula



wherein R is hydrogen

alkyl,  
hydroxyalkyl,  
alkoxyalkyl,  
trifluoromethyl,  
alkanoyloxyalkyl,  
alkanoylaminoalkyl,  
alkylthioalkyl,  
alkylsulfinylalkyl,  
alkylsulfonylalkyl,  
aminoalkyl,  
alkylaminoalkyl,  
dialkylaminoalkyl,  
N-alkylpiperazinoalkyl,  
N-phenylalkylpiperazinoalkyl,  
morpholinoalkyl,  
thiomorpholinoalkyl,  
piperidinoalkyl,  
pyrrolidinoalkyl,  
N-alkylalkylpiperidinoalkyl,  
pyridylalkyl,  
thienylalkyl,



- quinolinyllalkyl,  
thiazolyllalkyl,  
cycloalkyl,  
cycloalkylalkyl,  
5 phenyl,  
phenyl substituted by one to three substituents selected from the group  
consisting of:
- hydroxy,  
alkoxy,  
10 alkyl,  
alkylthio,  
alkylsulfinyl,  
alkylsulfonyl,  
amino,  
15 alkylamino,  
dialkylamino,  
halogen,  
cyano,  
nitro,  
20 trifluoromethyl or on adjacent carbon atoms by either a one to two  
carbon alkenylenedioxy group or a two to three carbon  
alkenyleneoxy group,  
phenylalkyl,  
phenylalkyl wherein phenyl is substituted by alkyl,  
25 alkoxy,  
halogen, or  
trifluoromethyl,  
heteroaryl,  
heteroaryl substituted by one to two substituents selected  
30 from the group consisting of:  
alkyl, or  
halogen,  
biphenyl,



-22-

biphenyl, substituted by alkyl,

alkoxy,

halogen,

trifluoromethyl, or

5

cyano,

biphenylalkyl or

biphenylalkyl wherein biphenyl is substituted by

alkyl,

alkoxy,

10

halogen,

trifluoromethyl, or

cyano;

D is zero or an integer of 1 to 3;

L is zero or an integer of 1 to 3;

15

R<sup>1a</sup> is the side chain of an unnaturally occurring amino acid. Unnaturally occurring amino acids are well-known in the art, e.g., Roberts et al., "Unusual Amino Acids in Peptide Synthesis," *The Peptides*, 1993;5:341-429, but are not naturally found in living organisms. The side chains of unnatural amino acids include, but are not limited to:

20

hydrogen,

-(CH<sub>2</sub>)<sub>n</sub>-naphthalimide wherein n is zero or an integer of 1 to 2,-(CH<sub>2</sub>)<sub>n</sub>-phthalimide wherein n is as defined above,-(CH<sub>2</sub>)<sub>n</sub>-aryl wherein n is as defined below,

alkyl,

25

substituted alkyl wherein the substituent is selected from the group consisting of:

SH,

OR<sup>5</sup> wherein R<sup>5</sup> is hydrogen, alkyl, phenyl, or benzyl,SR<sup>5</sup> wherein R<sup>5</sup> is as defined above,

30

halogen,






-23-

- $\text{NR}^5$   
 $|$   
 $\text{R}^{5a}$  wherein  $\text{R}^5$  and  $\text{R}^{5a}$  are either the same or different  
 and each is the same as defined above for  $\text{R}^5$ ,  
 $\text{CO}_2\text{H}$ ,  
 $\text{COR}^5$  wherein  $\text{R}^5$  is as defined above,  
 $\text{CHO}$ , or  
 $\text{CONR}^5$  wherein  $\text{R}^5$  and  $\text{R}^{5a}$  are either the same or  
 $|$   
 $\text{R}^{5a}$   
 different and each is the same as defined above for  
 $\text{R}^5$ ,  
 aryl,  
 $-(\text{CH}_2)_n$ -phenyl wherein  $n$  is as defined above,  
 alkenyl,  
 $(\text{CH}_2)_n$ -heteroaryl wherein  $n$  is as defined above,  
 heteroaryl,  
 heterocycle,  
 $-(\text{CH}_2)_m$ -NH-Z- $\text{R}^5$  wherein  $m$  is an integer of 1 to 6, Z is  
 $\begin{array}{cc} \text{O} & \text{S} \\ || & || \end{array}$   
 $-\text{C}-, -\text{C}-, \text{ or } -\text{SO}_2$  and  $\text{R}^5$  is as defined above,  
 $-(\text{CH}_2)_m$ -S-C(phenyl) $_3$  wherein  $m$  is as defined above,  
 $-(\text{CH}_2)_m$ -O-(CH $_2$ ) $L^a$ -phenyl wherein  $L^a$  is an integer of 1 to 6 and  
 $m$  is as defined above,  
 $\begin{array}{c} \text{O} \\ || \end{array}$   
 $-(\text{CH}_2)_m$ -C- $\text{R}^5$  wherein  $m$  and  $\text{R}^5$  are as defined above,  
 $-(\text{CH}_2)_m$ -NHSO $_2$ -aryl wherein  $m$  is as defined above,  
 $-(\text{CH}_2)_m$ -cycloalkyl wherein  $m$  is as defined above,  
 $\begin{array}{c} \text{O} \\ || \end{array}$   
 $-(\text{CH}_2)_m$ -C-aryl wherein  $m$  is as defined above,



-24-



Y is OR<sup>3</sup> wherein R<sup>3</sup> is hydrogen,

methyl,

10 ethyl, or

benzyl, or

NH-OR<sup>4</sup> wherein R<sup>4</sup> is hydrogen,

alkyl, or

benzyl;

15 with the proviso that R<sup>1a</sup> is not the side chain of a natural  $\alpha$  amino acid as defined above.

Additionally, side chains of natural  $\alpha$  amino acids and unnatural amino acids having the D or R configuration are included within this term.

20 The functional groups in the amino acid side chains can be protected. For example, carboxyl groups can be esterified, amino groups can be converted to amides or carbamates, hydroxyl groups can be converted to ethers or esters, and thiol groups can be converted to thioethers or thioesters.

"Halogen" is fluorine, chlorine, bromine, or iodine.

25 "Alkali metal" is a metal in Group IA of the periodic table and includes, for example, lithium, sodium, potassium, and the like.

The acyloxymethyl esters of compounds of Formula I can be prepared by methods known to one skilled in the art. For example, the corresponding carboxylic acids can be allowed to react first with a suitable base to give the carboxylate anion, followed by reaction with a carboxylic halomethyl ester, which



-25-

can be obtained from commercial suppliers or prepared by methods known to one skilled in the art, optionally in the presence of a suitable agent to activate the carboxylic halomethyl ester, which are known to one skilled in the art, to give the acyloxymethyl esters.

5           Some of the compounds of Formula I are capable of further forming both pharmaceutically acceptable acid addition and/or base salts. All of these forms are within the scope of the present invention.

          Pharmaceutically acceptable acid addition salts of the compounds of Formula I include salts derived from nontoxic inorganic acids such as  
10   hydrochloric, nitric, phosphoric, sulfuric, hydrobromic, hydriodic, hydrofluoric, phosphorous, and the like, as well as the salts derived from nontoxic organic acids, such as aliphatic mono- and dicarboxylic acids, phenyl-substituted alkanolic acids, hydroxy alkanolic acids, alkanedioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, etc. Such salts thus include sulfate, pyrosulfate, bisulfate,  
15   sulfite, bisulfite, nitrate, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, trifluoroacetate, propionate, caprylate, isobutyrate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, mandelate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, phthalate, benzenesulfonate,  
20   toluenesulfonate, phenylacetate, citrate, lactate, maleate, tartrate, methanesulfonate, and the like. Also contemplated are salts of amino acids such as arginate and the like and gluconate, galacturonate (see, for example, Berge S.M. et al., "Pharmaceutical Salts," *J. of Pharma. Sci.*, 1977;66:1).

          The acid addition salts of said basic compounds are prepared by contacting  
25   the free base form with a sufficient amount of the desired acid to produce the salt in the conventional manner. The free base form may be regenerated by contacting the salt form with a base and isolating the free base in the conventional manner. The free base forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are  
30   equivalent to their respective free base for purposes of the present invention.

          Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like.




Examples of suitable amines are N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge S.M. et al., "Pharmaceutical Salts," *J. of Pharma Sci.*, 1977;66:1).

5           The base addition salts of said acidic compounds are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in the conventional manner. The free acid forms differ from their respective salt  
10       forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention.

Certain of the compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms, including hydrated forms, are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention.

Certain of the compounds of the present invention possess one or more chiral centers and each center may exist in the R or S configuration. The present invention includes all diastereomeric, enantiomeric, and epimeric forms as well as the appropriate mixtures thereof. Additionally, the compounds of the present invention may exist as geometric isomers. The present invention includes all cis, trans, syn, anti, entgegen (E), and zusammen (Z) isomers as well as the appropriate mixtures thereof.

A preferred compound of Formula I is




wherein n, X, R, R<sup>1</sup>, and Y are as defined above.

Another preferred compound of Formula I is



-27-




I

wherein n is zero or an integer of 1; and

X is -O-, or

-CH<sub>2</sub>-.

5 Another preferred compound of Formula I is



I


wherein n is zero or an integer of 1;

X is -O-, or

-CH<sub>2</sub>-; and

10 R is hydrogen.

A more preferred compound of Formula I is



I

wherein n is zero or an integer of 1;


X is -O-, or

15 -CH<sub>2</sub>-;

R is hydrogen; and

Y is OH.

A most preferred compound of Formula I is



I

20 wherein n is zero or an integer of 1;



-28-

X is -O-, or

-CH<sub>2</sub>-;

R is hydrogen; and

Y is NHOH.

5           Particularly valuable in this embodiment of the invention is a compound selected from the group consisting of:

(S) 3-Methyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;

10           (S) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(S) 4-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;

(S) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-succinic acid;

15           (S) Phenyl-[(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)]-acetic acid;

(S) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-3-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-propionic acid;

(S) 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionic acid;

20           (S) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-4-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-butyric acid;

(S) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-5-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-pentanoic acid;

25           (S) 5-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid;

(S) 4-Phenylmethanesulfinyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;

(S) 4-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;

30           (S) 5-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid;



- (S) 6-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-hexanoic acid;
- (S) 7-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-heptanoic acid;
- 5 (S) 8-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-octanoic acid;
- (S) 4-Phenylsulfamoyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- (S) 4-Phenylmethanesulfonyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- 10 (S) 4-Benzylsulfanyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- (S) 3-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionic acid;
- 15 (S) 4-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- (S) 5-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid;
- (S) 6-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-hexanoic acid;
- 20 (S) 7-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-heptanoic acid;
- (S) 8-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-octanoic acid;
- 25 (S) 2-(2,3-Dihydro-1H-8-oxa-cyclopenta[a]indene-6-sulfonylamino)-3-methyl-butyric acid;
- (S) 3-Methyl-2-(6,7,8,9-tetrahydro-5H-10-oxa-benzo[a]azulene-2-sulfonylamino)-butyric acid;
- (S) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;
- 30 (S) N-Hydroxy-3-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-succinamic acid;



(S) N-Hydroxy-2-phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-acetamide;

(S) N-Hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-3-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-propionamide;

5 (S) 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-N-hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionamide;

(S) N-Hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-4-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-butyramide;

10 (S) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-5-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-pentanoic acid hydroxyamide;

(S) 5-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid hydroxyamide;

(S) N-Hydroxy-4-phenylmethanesulfinyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

15 (S) 4-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-N-hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(S) 5-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid hydroxyamide;

20 (S) 6-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-hexanoic acid hydroxyamide;

(S) 7-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-heptanoic acid hydroxyamide;

(S) 8-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-octanoic acid hydroxyamide;

25 (S) 4-Benzylsulfonyl-N-hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(S) N-Hydroxy-4-phenylsulfamoyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

30 (S) N-Hydroxy-4-phenylmethanesulfonyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(S) N-Hydroxy-3-(1H-indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionamide;



- (S) N-Hydroxy-4-(1H-indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;
- (S) 5-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid hydroxyamide;
- 5 (S) 6-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-hexanoic acid hydroxyamide;
- (S) 7-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-heptanoic acid hydroxyamide;
- (S) 8-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-octanoic acid hydroxyamide;
- 10 (S) 2-(2,3-Dihydro-1H-8-oxa-cyclopenta[a]indene-6-sulfonylamino)-N-hydroxy-3-methyl-butylamide;
- (S) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-5H-10-oxa-benzo[a]azulene-2-sulfonylamino)-butyramide;
- 15 (S) 3-Methyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyric acid;
- (S) 3-Methyl-2-(9-methyl-6,7,8,9-tetrahydro-5H-carbazole-2-sulfonylamino)-butyric acid;
- (S) 4-Phenyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyric acid;
- 20 (S) 4-Phenyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyric acid;
- (S) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyramide;
- 25 (S) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyramide;
- (S) N-Hydroxy-3-methyl-2-(9-methyl-6,7,8,9-tetrahydro-5H-carbazole-2-sulfonylamino)-butyramide;
- (S) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyramide;
- 30 (S) 3-Methyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyric acid;



- (S) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyramide;
- (R) 3-Methyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- 5 (R) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;
- (R) 4-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- (R) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-succinic acid;
- 10 (R) Phenyl-[(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)]-acetic acid;
- (R) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-3-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-propionic acid;
- (R) 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionic acid;
- 15 (R) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-4-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-butyric acid;
- (R) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-5-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-pentanoic acid;
- 20 (R) 5-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid;
- (R) 4-Phenylmethanesulfinyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- (R) 4-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- 25 (R) 5-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid;
- (R) 6-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-hexanoic acid;
- 30 (R) 7-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-heptanoic acid;
- (R) 8-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-octanoic acid;



- (R) 4-Phenylsulfamoyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- (R) 4-Phenylmethanesulfonyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- 5 (R) 4-Benzylsulfanyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- (R) 3-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionic acid;
- (R) 4-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- 10 (R) 5-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid;
- (R) 6-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-hexanoic acid;
- 15 (R) 7-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-heptanoic acid;
- (R) 8-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-octanoic acid;
- (R) 2-(2,3-Dihydro-1H-8-oxa-cyclopenta[a]indene-6-sulfonylamino)-3-methyl-butyl-butyric acid;
- 20 (R) 3-Methyl-2-(6,7,8,9-tetrahydro-5H-10-oxa-benzo[a]azulene-2-sulfonylamino)-butyric acid;
- (R) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;
- 25 (R) N-Hydroxy-3-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-succinamic acid;
- (R) N-Hydroxy-2-phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-acetamide;
- (R) N-Hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-3-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-propionamide;
- 30 (R) 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-N-hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionamide;



(R) N-Hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-4-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-butyramide;

(R) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-5-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-pentanoic acid hydroxyamide;

5 (R) 5-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid hydroxyamide;

(R) N-Hydroxy-4-phenylmethanesulfinyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

10 (R) 4-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-N-hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(R) 5-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid hydroxyamide;

(R) 6-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-hexanoic acid hydroxyamide;

15 (R) 7-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-heptanoic acid hydroxyamide;

(R) 8-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-octanoic acid hydroxyamide;

20 (R) 4-Benzylsulfonyl-N-hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(R) N-Hydroxy-4-phenylsulfamoyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(R) N-Hydroxy-4-phenylmethanesulfonyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

25 (R) N-Hydroxy-3-(1H-indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionamide;

(R) N-Hydroxy-4-(1H-indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

30 (R) 5-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid hydroxyamide;

(R) 6-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-hexanoic acid hydroxyamide;



-35-

(R) 7-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-heptanoic acid hydroxyamide;

(R) 8-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-octanoic acid hydroxyamide;

5 (R) 2-(2,3-Dihydro-1H-8-oxa-cyclopenta[a]indene-6-sulfonylamino)-N-hydroxy-3-methyl-butyramide;

(R) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-5H-10-oxa-benzo[a]azulene-2-sulfonylamino)-butyramide;

10 (R) 3-Methyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyric acid;

(R) 3-Methyl-2-(9-methyl-6,7,8,9-tetrahydro-5H-carbazole-2-sulfonylamino)-butyric acid;

(R) 4-Phenyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyric acid;

15 (R) 4-Phenyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyric acid;

(R) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyramide;

20 (R) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyramide;

(R) N-Hydroxy-3-methyl-2-(9-methyl-6,7,8,9-tetrahydro-5H-carbazole-2-sulfonylamino)-butyramide;

(R) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyramide;

25 (R) 3-Methyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyric acid; and

(R) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyramide;

and corresponding isomers thereof; or a pharmaceutically acceptable salt thereof.

30 The compounds of Formula I are valuable inhibitors of a number of different matrix metalloproteinases. It has been shown previously that inhibitors of matrix metalloproteinases have efficacy in models of disease states like arthritis and metastasis that depend on modification of the extracellular matrix.



In vitro experiments were carried out which demonstrate the efficacy of compounds of Formula I as potent and specific inhibitors of a variety of matrix metalloproteinases. Experiments were carried out with the full-length and catalytic domains of the proteinases. Table 1 shows the activity of Examples 1-4 versus

5 MMP-1FL (collagenase-1 full length enzyme), MMP-2CD (gelatinase A catalytic domain), MMP-2FL (gelatinase A full length enzyme), MMP-3CD (stromelysin-1 catalytic domain), MMP-7FL (matrilysin full length enzyme), MMP-9-FL (gelatinase B full length enzyme), MMP-13CD (collagenase-3 catalytic domain), and MMP-14CD (membrane-type MMP-1). IC<sub>50</sub> values were determined using a

10 thiopeptolide substrate, Ac-Pro-Leu-Gly-thioester-Leu-Leu-Gly-OEt (Ye Q.-Z., Johnson L.L., Hupe D.J., and Baragi V., "Purification and Characterization of the Human Stromelysin Catalytic Domain Expressed in *Escherichia coli*," *Biochemistry*, 1992;31:11231-11235; Ye Q.-Z., Johnson L.L., Yu A.E., and Hupe D., "Reconstructed 19 kDa catalytic domain of gelatinase A is an active

15 proteinase," *Biochemistry*, 1995;34:4702-4708.) MMP-13CD was expressed from a synthetic gene and purified from *Escherichia coli* cell culture according to a previously described method (Ye Q.-Z., Johnson L.L., and Baragi V., "Gene synthesis and expression in *E. coli* for PUMP, a human matrix metalloproteinase," *Biochemical and Biophysical Research Communications*, 1992;186:143-149).



TABLE 1. Biological Activity of Compounds of Formula I

Example	IC <sub>50</sub> (μM)									
	MMP-1FL	MMP-2CD	MMP-2FL	MMP-3CD	MMP-7FL	MMP-9FL	MMP-13CD	MMP-14CD		
1	6.7	0.037	0.335	0.046	82	30	3.45	0.38		
2	6.05	0.034	0.43	0.0245	7.5	100	2.3	0.155		
3	16	0.0167	0.45	0.0056	3.2	100	0.97	0.048		
4	25	0.018	0.15	0.0215	2.3	28	1.345	0.064		



The following list contains abbreviations and acronyms used within the schemes and text:


GBM	Glomerular basement membrane
ECM	Extracellular matrix
CNS	Central nervous system
CH <sub>2</sub> Cl <sub>2</sub>	Dichloromethane
EAE	Experimental autoimmune encephalomyelitis
MMP	Matrix metalloproteinase
TIMPs	Tissue inhibitors of matrix metalloproteinases
VSMC	Vascular smooth muscle cell
TFA	Trifluoroacetic acid
IC <sub>50</sub>	Concentration of compound required to inhibit 50% of enzyme activity
HCl	Hydrogen chloride
THF	Tetrahydrofuran
Pd	Palladium
Na	Sodium
NaH	Sodium hydride
LiOH	Lithium hydroxide
LiCl	Lithium Chloride
H <sub>2</sub> O	Water
H <sub>2</sub>	Hydrogen
CDI	1,1'-Carbonyldiimidazole
Hv	light
SO <sub>3</sub> -DMF	Sulfur trioxide dimethyl formamide
SOCl <sub>2</sub>	Thionyl Chloride
t-Bu	tertiary butyl
BOC	tertiary butoxycarbonyl
LDA	Lithium diisopropylamide
MeOH	Methanol
DMF	Dimethylformamide



-39-

p-TsOH(p-TSA)	para-Toluenesulfonic acid
CHCl <sub>3</sub>	Chloroform
CDCl <sub>3</sub>	Deuterated chloroform
E	Entgegen
Z	Zusammen
H <sub>2</sub> NOBz	O-Benzyl hydroxylamine
TEA	Triethylamine
CH <sub>3</sub> CN	Acetonitrile
DBU	1,8-Diazabicyclo[5.4.0]undec-7-ene
DCC	Dicyclohexylcarbodiimide
PPA	Polyphosphoric acid
BaSO <sub>4</sub>	Barium sulfate
DMSO-d <sub>6</sub>	Deuterated dimethylsulfoxide
MgSO <sub>4</sub>	Magnesium sulfate
<sup>1</sup> H-NMR	Proton nuclear magnetic resonance
PPM	Parts per million
MS	Mass spectrum

Tricyclic aryl and tricyclic heteroaryl starting materials of formula (3)



wherein n is zero or an integer of 1 or 2; and

X is -O-,

-S-,

-N-, wherein R<sup>2</sup> is hydrogen,

|  
R<sup>2</sup>

alkyl,

acyl, or

benzyl,



are either obtained from commercial sources ( $X = N-R^2$  wherein  $R^2$  is as defined above) or prepared using methods known in the art, e.g., Bachelet J.P. and Caubere P., *J. Org. Chem.*, 1982;47:234-238; Ebel F., *Helv. Chim. Acta*, 1929;12:3-16; Vanrysselberghe V. et al., *Ind. Eng. Chem. Res.*, 1996;35:3311-3318; Derouane D. et al., *J. Chem. Soc., Chem. Commun.*, 1995;10:993-994; Miki Y. and Sugimoto Y., *Seikiyu Gakkaishi*, 1994;37:386-394; Miki Y. et al., *Seikiyu Gakkaishi*, 1992;35:332-338; Rankel L.A., *Fuel Sci. Technol. Int.*, 1991;9:1435-1447; Siskin M. et al., *Energy Fuels*, 1990;4:482-488; Sundaram K.M. et al., *Chem. Eng. Commun.*, 1988;71:53-71; Francisco M.A. et al., *J. Org. Chem.*, 1988;53:596-600; Nagai M. et al., *J. Catal.*, 1986;97:52-58; Miyake M. et al., *Bull. Chem. Soc. Japan*, 1979;52:559-563; Ando W. et al., *J. Chem. Soc. Chem. Commun.*, 1975;17:704-705; Fraser P.S. et al., *J. Org. Chem.*, 1974;39:2509-2513; Cagniant P. et al., *Bull. Soc. Chim. Fr.*, 1969;2:607-612; and Cagniant D. et al., *Bull. Soc. Chim. Fr.*, 1969;2:601-606; United States Patents 5,721,185, 5,670,680; International Published Patent Application WO 95/27717; Smith W. et al., *J. Org. Chem.*, 1990;55:5301-5302; Mejer S., *Pol. of Chem.*, 1979;53:2385-2388; Canonne P. et al., *J. Org. Chem.*, 1980;45:1828-1835; Parham W.E., *Synthesis*, 1976;116-117; Japanese Patent Application JP 08191063 A2; Parhan W.E., *J. Org. Chem.*, 1969;34:1899-1904; McClure K.F. et al., *Bioorg. Med. Chem. Lett.*, 1998;8:143-146.

The synthesis of starting materials for a compound of Formula I wherein X is -O- is shown in Scheme 1. Thus, a compound of formula (1) wherein n is zero or an integer of 1 or 2 is reacted with phenol in the presence of sodium, or sodium hydride and the like in the presence of a solvent such as benzene, tetrahydrofuran and the like to afford a compound of formula (2). Cyclization of a compound of formula (2) in the presence of an acid such as, for example, polyphosphoric acid, para-toluenesulfonic acid and the like in the presence of a solvent such as benzene and the like affords a compound of formula (3).

The synthesis of compounds of Formula Ia, Ib, Ic, and Id are shown in Scheme 2. Thus, a compound of formula (3) wherein n is as defined above is sulfonated using a sulfonating reagent such as, for example,  $SO_3$ -DMF, and the like by refluxing in a solvent such as, for example, dichloroethane and the like to



-41-

afford a compound of formula (4) wherein n is as defined above. A compound of formula (4) is chlorinated with a chlorinating reagent such as, for example, thionyl chloride and the like at about room temperature to afford a compound of formula (5) wherein n is as defined above. A compound of formula (5) is reacted with an amino acid of formula (6) wherein R<sup>1</sup> is hydrogen, a side chain of a natural amino acid or a side chain of an unnatural amino acid in the presence of a base such as, for example, triethylamine and the like in a solvent such as, for example, tetrahydrofuran/water and the like at about room temperature to afford a compound of Formula Ia wherein n and R<sup>1</sup> are as defined above. Alternatively, a compound of formula (5) is reacted with a C-protected amino acid of formula (7) wherein R<sup>1</sup> is as defined above in the presence of a base such as, for example, triethylamine and the like in the presence of a solvent such as, for example, dichloromethane and the like to afford a compound of formula (8) wherein n and R<sup>1</sup> are as defined above. A compound of formula (8) can be deprotected in the presence of an acid such as, for example, trifluoroacetic acid and the like, and a solvent such as, for example, dichloromethane and the like at about room temperature to afford a compound of Formula Ia. Coupling the acid chloride of Formula Ia with O-benzyl hydroxylamine in a solvent such as, for example, tetrahydrofuran and the like at about -10°C to about 40°C to afford a compound of Formula Ic wherein n and R<sup>1</sup> are as defined above. Reaction of a compound of Formula Ic with hydrogen gas in the presence of a catalyst such as, for example, palladium on barium sulfate and the like in a solvent such as, for example, methanol, tetrahydrofuran and the like to afford a compound of Formula Id wherein n and R<sup>1</sup> are as defined above. Reaction of a compound of formula (8) with a compound of formula

R-Hal

wherein R is hydrogen,  
alkyl,  
hydroxyalkyl,  
alkoxyalkyl,  
trifluoromethyl,  
alkanoyloxyalkyl,



- alkanoylaminoalkyl,  
alkylthioalkyl,  
alkylsulfinylalkyl,  
alkylsulfonylalkyl,  
5 aminoalkyl,  
alkylaminoalkyl,  
dialkylaminoalkyl,  
N-alkylpiperazinoalkyl,  
N-phenylalkylpiperazinoalkyl,  
10 morpholinoalkyl,  
thiomorpholinoalkyl,  
piperidinoalkyl,  
pyrrolidinoalkyl,  
N-alkylalkylpiperidinoalkyl,  
15 pyridylalkyl,  
thienylalkyl,  
quinolinyllalkyl,  
thiazolylalkyl,  
cycloalkyl,  
20 cycloalkylalkyl,  
phenyl,  
phenyl substituted by one to three substituents selected from the group  
consisting of:
- 25 hydroxy,  
alkoxy,  
alkyl,  
alkylthio,  
alkylsulfinyl,  
alkylsulfonyl,  
30 amino,  
alkylamino,  
dialkylamino,  
halogen,



-43-

cyano,

nitro,

trifluoromethyl or on adjacent carbon atoms by either a one to two

carbon alkenylenedioxy group or a two to three carbon

5 alkenyleneoxy group,

phenylalkyl,

phenylalkyl wherein phenyl is substituted by alkyl,

alkoxy,

halogen, or

10 trifluoromethyl,

heteroaryl,

heteroaryl substituted by one to two substituents selected from the  
group consisting of:

alkyl, or

15 halogen,

biphenyl,

biphenyl, substituted by alkyl,

alkoxy,

halogen,

20 trifluoromethyl, or

cyano,

biphenylalkyl or

biphenylalkyl wherein biphenyl is substituted by

alkyl,

25 alkoxy,

halogen,

trifluoromethyl, or

cyano;

30 and Hal is chlorine, bromine, or iodine in the presence of a base such as, for  
example, DBU and the like in a solvent such as, for example, acetonitrile and the  
like to afford a compound of formula (9) wherein n, R, and R<sup>1</sup> are as defined  
above. Reaction of a compound of formula (9) with an acid such as, for example,



trifluoroacetic acid and the like in the presence of a solvent such as, for example, dichloromethane affords a compound of Formula Ib wherein n, R, and R<sup>1</sup> are as defined above.


The previous methodology can be applied to both natural and unnatural  
 5  $\alpha$ -amino acids of formulas (6) and (7) which are readily available from commercial sources or can be prepared by methods known in the art. Alternatively, natural and unnatural  $\alpha$ -amino acids of formulas (6) and (7) can be prepared as shown in Scheme 3. Thus employing the method of Evans D.A. et al.,  
 10 *J. Amer. Chem. Soc.*, 1982;104:1737-1739, N-Boc-glycine of formula (10) is coupled with the chiral sodium salt of benzyl oxazolidine of formula (11) in the presence of a coupling reagent such as, for example, carbonyldiimidazole in a solvent such as, for example, tetrahydrofuran and the like at about -10°C to about room temperature to afford the compound of formula (12). The enolate of the compound of formula (12) is formed by reaction with lithium diisopropylamide  
 15 and subsequently alkylated with a compound of formula



wherein R<sup>1</sup> and Hal are as defined above to afford a compound of formula (13) as a mixture of diastereomers. The diastereomers are separated using  
 20 chromatography using an absorbent such as, for example, silica gel and the like to afford pure diastereomers. The pure diastereomers are treated with gaseous hydrogen chloride in a solvent such as dichloromethane and the like at about room temperature to afford a compound of formula (14) wherein R<sup>1</sup> is as defined above. Reaction of a compound of formula (14) with a compound of formula



25 wherein Ar is



wherein

X is -O-,



-45-

-S(O)<sub>p</sub>- wherein p is zero or an integer of 1 or 2,

-N-, wherein R<sup>2</sup> is hydrogen,

|                      alkyl,

R<sup>2</sup>                      acyl, or

5                              benzyl,

-CH<sub>2</sub>-, or

-C-,

||

O

10 and n is as defined above (prepared by methodology described previously for preparing a compound of formula (5) from a compound of formula (3)) in the presence of a base such as, for example, triethylamine and the like, in a solvent mixture such as, for example, tetrahydrofuran and water at about 10°C to about room temperature to afford a compound of formula (15) wherein R<sup>1</sup> is as defined  
15 above. The oxazolidone chiral auxiliary is removed by hydrolysis with a base such as, for example, lithium hydroxide and the like, in a solvent mixture such as, for example, dioxane/water at about room temperature to afford a compound of Formula 1e wherein R<sup>1</sup> is as defined above.

20 Scheme 4 shows the preparation of a compound of Formula 1f using the methodology of Myers A.G. et al., *Tetrahedron Lett.*, 1995;36:4555-4558. Thus, the enantiomeric pseudoephedrine glycinamide of formula (16) or its enantiomer is added to a slurry of lithium chloride and lithium diisopropylamide in a solvent such as, for example, tetrahydrofuran at about -78°C to afford an O,N-dianion which is warmed to about 0°C and treated with a compound of formula

25  $R^1\text{-Hal}$

wherein R<sup>1</sup> and Hal are as defined above to afford a compound of formula (17)

wherein R<sup>1</sup> is as defined above having a high degree of diastereoselectivity

(>99% de). A compound of formula (17) is treated with a compound of formula

$\text{Ar SO}_2\text{Cl}$

30 wherein Ar is as defined above in the presence of a base such as, for example, triethylamine and the like in a solvent such as, for example, tetrahydrofuran/water at about 10°C to about room temperature to afford a compound of formula (18)



-46-


wherein Ar and R<sup>1</sup> are as defined above. The chiral auxiliary is then removed by hydrolysis with a base such as, for example, aqueous sodium hydroxide or water/methanol mixtures at about reflux to afford a compound of Formula If wherein Ar and R<sup>1</sup> are as defined above. Alternatively, the hydrolysis can be  
5 carried out by refluxing an aqueous solution of a compound of formula (18) without additional base being added.

Compounds of formula R-Hal or R<sup>1</sup>-Hal are either commercially available or can be obtained by methods known in the art.



-47-


Scheme 1





-48-


Scheme 2





-49-


Scheme 3





-50-

Scheme 4



The compounds of the present invention can be prepared and administered in a wide variety of oral and parenteral dosage forms. Thus, the compounds of the present invention can be administered by injection, that is, intravenously, intramuscularly, intracutaneously, subcutaneously, intraduodenally, or intraperitoneally. Also, the compounds of the present invention can be administered by inhalation, for example, intranasally. Additionally, the compounds of the present invention can be administered transdermally. It will be obvious to those skilled in the art that the following dosage forms may comprise as the active component, either a compound of Formula I or a corresponding pharmaceutically acceptable salt of a compound of Formula I.

For preparing pharmaceutical compositions from the compounds of the present invention, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substances which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.



In powders, the carrier is a finely divided solid which is in a mixture with the finely divided active component.

In tablets, the active component is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

The powders and tablets preferably contain from 5 or 10 to about 70% of the active compound. Suitable carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as a carrier providing a capsule in which the active component, with or without other carriers, is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid dosage forms suitable for oral administration.

For preparing suppositories, a low melting wax, such as a mixture of fatty acid glycerides or cocoa butter, is first melted and the active component is dispersed homogeneously therein, as by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and thereby to solidify.

Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water propylene glycol solutions. For parenteral injection, liquid preparations can be formulated in solution in aqueous polyethylene glycol solution.

Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizing, and thickening agents as desired.

Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents.

Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for oral administration.



-52-

Such liquid forms include solutions, suspensions, and emulsions. These preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

5. The pharmaceutical preparation is preferably in unit dosage form. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit  
10 dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

- The quantity of active component in a unit dose preparation may be varied or adjusted from 1 mg to 1000 mg, preferably 10 mg to 100 mg according to the particular application and the potency of the active component. The composition  
15 can, if desired, also contain other compatible therapeutic agents.

- In therapeutic use as agents for the treatment of multiple sclerosis, atherosclerotic plaque rupture, aortic aneurism, heart failure, left ventricular dilation, restenosis, periodontal disease, corneal ulceration, treatment of burns, decubital ulcers, wound healing, cancer, inflammation, pain, arthritis,  
20 osteoporosis, renal disease, or other autoimmune or inflammatory disorders dependent upon tissue invasion by leukocytes, or other activated migrating cells, acute and chronic neurodegenerative disorders including stroke, head trauma, spinal cord injury, Alzheimer's disease, amyotrophic lateral sclerosis, cerebral amyloid angiopathy, AIDS, Parkinson's disease, Huntington's disease, prion  
25 diseases, myasthenia gravis, and Duchenne's muscular dystrophy, the compounds utilized in the pharmaceutical methods of the invention are administered at the initial dosage of about 1 mg to about 100 mg per kilogram daily. A daily dose range of about 25 mg to about 75 mg per kilogram is preferred. The dosages, however, may be varied depending upon the requirements of the patient, the  
30 severity of the condition being treated, and the compound being employed. Determination of the proper dosage for a particular situation is within the skill of the art. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small



increments until the optimum effect under the circumstance is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired.

5 The following nonlimiting examples illustrate the inventors' preferred methods for preparing the compounds of the invention.

#### EXAMPLE 1

##### (R) 3-Methyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid

##### Step (a) Preparation of 6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonic acid

To a solution of tetrahydrodibenzofuran (4 g, 0.023 mol) in dichloroethane  
10 (50 mL) was added, in one portion, sulfur dioxide-DMF complex (6 g, 0.039 mol). The reaction mixture was refluxed for 14 hours, cooled, and concentrated in vacuo. The resulting crude liquid was dissolved in warm diethyl ether/ethanol yielding a precipitate on cooling. The solid was collected by filtration, washed with diethyl ether, and dried in vacuo to give the title compound as a pink solid  
15 (2.3 g, 40%).  
 $^1\text{H}$ NMR ( $\text{CDCl}_3$ )  $\delta$  7.9 (s, 1H), 7.7 (d, 1H), 7.4 (d, 1H), 2.8 (m, 2H), 2.6 (m, 2H), 2.0-1.8 (m, 4H) ppm.

##### Step (b) Preparation of 6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonyl chloride

6,7,8,9-Tetrahydrodibenzofuran-3-sulfonic acid (2.1 g, 8.3 mmol) was  
20 suspended in thionyl chloride (25 mL) and stirred at room temperature for 6 hours. The solution was concentrated in vacuo, and the resulting liquid was taken up in ethyl acetate, washed with water, brine, and dried over magnesium sulfate. The solvent was concentrated to dryness, and the crude product was triturated with hexane and collected by filtration to give the sulfonyl chloride as a tan solid  
25 (1.3 g, 58%).  
 $^1\text{H}$ NMR ( $\text{CDCl}_3$ )  $\delta$  8.0 (s, 1H), 7.8 (d, 1H), 7.5 (d, 1H), 2.8 (m, 2H), 2.6 (m, 2H), 1.9-1.7 (m, 4H) ppm.



-54-

Step (c) Preparation of (R) 3-Methyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid, *tert*-butyl ester

To a solution of (D)-valine, *tert*-butyl ester (0.13 g, 0.74 mmol) and triethylamine (0.075 g, 0.74 mmol) in tetrahydrofuran/water (5 mL, 1:1) was added in one portion at room temperature 6,7,8,9-tetrahydro-dibenzofuran-3-sulfonyl chloride (0.20 g, 0.74 mmol). The reaction mixture was stirred at room temperature for 14 hours, followed by the addition of aqueous HCl (1 M, 5 mL) and ethyl acetate (10 mL). The organic phase was separated and washed with brine, dried (MgSO<sub>4</sub>), and concentrated to dryness to give the title compound as a white solid (0.24 g, 80%).

<sup>1</sup>HNMR (CDCl<sub>3</sub>) δ 7.9 (s, 1H), 7.7 (d, 1H), 7.5 (d, 1H), 5.1 (d, 1H), 3.6 (dd, 1H), 2.8 (m, 2H), 2.6 (m, 2H), 2.0 (m, 1H), 1.9-1.8 (m, 4H), 1.1 (s, 9H), 1.0 (d, 3H), 0.8 (d, 3H) ppm.

Step (d) Preparation of (R) 3-Methyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid

To a solution of anisole (0.062 g, 0.57 mmol) in trifluoroacetic acid (3 mL) stirred at room temperature was added (R) 3-methyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid, *tert*-butyl ester (0.23 g, 0.56 mmol). Hydrolysis of the ester was completed in 4 hours, at which time the acidic solution was poured over ice, and the resulting solid was collected by filtration. The filter cake was dried in vacuo, and the solid was recrystallized from hexane/ethyl acetate to yield a cream-colored powder (0.12 g, 71%); mp 167-169°C.

<sup>1</sup>HNMR (CDCl<sub>3</sub>) δ 7.9 (s, 1H), 7.7 (d, 1H), 7.4 (d, 1H), 5.4 (d, 1H), 3.7 (dd, 1H), 2.7 (m, 2H), 2.5 (m, 2H), 2.1 (m, 1H), 1.9-1.7 (m, 4H), 0.9 (d, 3H), 0.8 (d, 3H) ppm.

In a manner similar to that described in Example 1, the following compounds were prepared:



-55-

## EXAMPLE 2

(S) 3-Methyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;

mp 162-165°C.

<sup>1</sup>HNMR (CDCl<sub>3</sub>) δ 7.9 (s, 1H), 7.7 (d, 1H), 7.5 (d, 1H), 5.1 (d, 1H), 3.8 (dd, 1H),

5 2.8 (m, 2H), 2.6 (m, 2H), 2.1 (m, 1H), 2.0-1.8 (m, 4H), 0.9 (d, 3H),

0.8 (d, 3H) ppm.

## EXAMPLE 3

(S) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-succinic acid;

mp 176-179°C.

10 <sup>1</sup>HNMR (CDCl<sub>3</sub>/DMSO-d<sub>6</sub>) δ 7.8 (s, 1H), 7.6 (d, 1H), 7.4 (d, 1H), 5.9 (d, 1H),

3.9 (m, 1H), 2.9-2.5 (m, 7H), 1.9-1.7 (m, 4H) ppm.

In a manner similar to that described in Example 1 but replacing D-valine, *tert*-butyl ester with L-homophenylalanine, methyl ester and utilizing basic hydrolysis of the ester moiety, the following compound was prepared:

15

## EXAMPLE 4

(S) 4-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;

mp 167-169°C.

<sup>1</sup>HNMR (CDCl<sub>3</sub>) δ 7.9 (s, 1H), 7.7 (d, 1H), 7.5 (d, 1H), 7.2 (m, 3H), 7.1 (d, 2H),

5.2 (d, 1H), 4.0 (m, 1H).

20

Using the procedure of Example 1, the following compounds of Formula I are prepared:

## EXAMPLE 5

(S)-Phenyl-[(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)]-acetic acid

## EXAMPLE 6

25

(S) 5-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid



-56-

## EXAMPLE 7

(S) 2-(2,3-Dihydro-1H-8-oxa-cyclopenta[a]indene-6-sulfonylamino)-3-methyl-  
butyric acid

## EXAMPLE 8

5 (S) 3-Methyl-2-(6,7,8,9-tetrahydro-5H-10-oxa-benzo[a]azulene-2-sulfonylamino)-  
butyric acid

## EXAMPLE 9

(S) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-  
butyramide

10

## EXAMPLE 10

(S) N-Hydroxy-3-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-succinamic  
acid

## EXAMPLE 11

(R) 4-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid

15

## EXAMPLE 12

(R) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-succinic acid

## EXAMPLE 13

(R) 2-(2,3-Dihydro-1H-8-oxa-cyclopenta[a]indene-6-sulfonylamino)-3-methyl-  
butyric acid

20

## EXAMPLE 14

(R) 3-Methyl-2-(6,7,8,9-tetrahydro-5H-10-oxa-benzo[a]azulene-2-  
sulfonylamino)-butyric acid

## EXAMPLE 15

25

(S) 3-Methyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyric  
acid



-57-

## EXAMPLE 16

(S) 3-Methyl-2-(9-methyl-6,7,8,9-tetrahydro-5H-carbazole-2-sulfonylamino)-butyric acid

## EXAMPLE 17

5 (S) 4-Phenyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyric acid;

## EXAMPLE 18

(S) 3-Methyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyric acid

GENERAL PROCEDURE FOR PREPARING HYDROXAMIC ACIDS OF  
FORMULA I (Y = NHOH)

10 Step (a) Preparation of O-Benzylhydroxylamine derivative

To a solution of the acid chloride of a carboxylic acid of Formula I (Y = OH) in tetrahydrofuran is added two equivalents of O-benzylhydroxylamine (the acid chloride is prepared from the corresponding acid and thionyl chloride or oxalyl chloride). The mixture is stirred at room temperature for 24 hours. The  
15 reaction mixture is diluted with 1 M hydrochloric acid and ethyl acetate. The layers are separated and the ethyl acetate solution is washed with water and dried over magnesium sulfate. After filtration, the crude solid is triturated with hexane, and the resulting solid is collected by filtration.

Step (b) Preparation of Hydroxamic Acid of Formula I (Y = NHOH)

20 To a solution of the product from Step (a) in methanol/tetrahydrofuran is added palladium on barium sulfate. The solution is exposed to hydrogen gas either at atmospheric pressure under a balloon or at 50 pounds per square inch (psi) in a Parr apparatus. After hydrogen uptake is complete, the mixture is filtered through celite and concentrated in vacuo to afford the hydroxamic acid.

25 Using the general procedure for preparing hydroxamic acids described above, the following hydroxamic acids of Formula I are prepared.



-58-

## EXAMPLE 19

(S) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-  
butyramide

## EXAMPLE 20

5 (S) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-  
butyramide

## EXAMPLE 21

(S) N-Hydroxy-2-phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-  
acetamide

10

## EXAMPLE 22

(S) N-Hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-3-(3,4,4-  
trimethyl-2,5-dioxo-imidazolidin-1-yl)-propionamide

## EXAMPLE 23

15 (S) 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-N-hydroxy-2-(6,7,8,9-tetrahydro-  
dibenzofuran-3-sulfonylamino)-propionamide

## EXAMPLE 24

(S) 5-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid  
hydroxyamide

## EXAMPLE 25

20 (S) N-Hydroxy-3-(1H-indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-  
3-sulfonylamino)-propionamide

## EXAMPLE 26

(S) 2-(2,3-Dihydro-1H-8-oxa-cyclopenta[a]indene-6-sulfonylamino)-N-hydroxy-  
3-methyl-butyramide



## EXAMPLE 27

(S) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-5H-10-oxa-benzo[a]azulene-2-sulfonylamino)-butyramide

## EXAMPLE 28

5 (S) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyramide

## EXAMPLE 29

(S) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyramide

10

## EXAMPLE 30

(R) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide

## EXAMPLE 31

15

(R) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide

## EXAMPLE 32

(R) N-Hydroxy-2-phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-acetamide

## EXAMPLE 33

20

(R) N-Hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-3-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-propionamide

## EXAMPLE 34

(R) 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-N-hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionamide



-60-

## EXAMPLE 35

(R) 5-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid hydroxyamide

## EXAMPLE 36

5 (R) N-Hydroxy-3-(1H-indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionamide

## EXAMPLE 37

(R) 2-(2,3-Dihydro-1H-8-oxa-cyclopenta[a]indene-6-sulfonylamino)-N-hydroxy-3-methyl-butyramide

10

## EXAMPLE 38

(R) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-5H-10-oxa-benzo[a]azulene-2-sulfonylamino)-butyramide

## EXAMPLE 39

15 (R) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyramide

## EXAMPLE 40


(R) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyramide



-61-

## CLAIMS

1. A compound of Formula I



wherein n is zero or an integer of 1 or 2;

X is -O-,

-S(O)<sub>p</sub> wherein p is zero or an integer of 1 or 2,

-N- wherein R<sup>2</sup> is hydrogen,

alkyl,  
|  
R<sup>2</sup> acyl, or  
benzyl,

-CH<sub>2</sub>-, or

-C-;  
||  
O

R is hydrogen,

alkyl,

hydroxyalkyl,

alkoxyalkyl,

trifluoromethyl,

alkanoyloxyalkyl,

alkanoylaminoalkyl,

alkylthioalkyl,

alkylsulfinylalkyl,

alkylsulfonylalkyl,

aminoalkyl,

alkylaminoalkyl,

dialkylaminoalkyl,

N-alkylpiperazinoalkyl,



-62-

5 N-phenylalkylpiperazinoalkyl,  
morpholinoalkyl,  
thiomorpholinoalkyl,  
piperidinoalkyl,  
pyrrolidinoalkyl,  
N-alkylalkylpiperidinoalkyl,  
pyridylalkyl,  
thienylalkyl,  
quinolinyalkyl,  
10 thiazolylalkyl,  
cycloalkyl,  
cycloalkylalkyl,  
phenyl,  
phenyl substituted by one to three substituents selected from the  
15 group consisting of:  
hydroxy,  
alkoxy,  
alkyl,  
alkylthio,  
20 alkylsulfinyl,  
alkylsulfonyl,  
amino,  
alkylamino,  
dialkylamino,  
25 halogen,  
cyano,  
nitro,  
trifluoromethyl or on adjacent carbon atoms by either a one  
to two carbon alkenylenedioxy group or a two to  
30 three carbon alkenyleneoxy group,  
phenylalkyl,  
phenylalkyl wherein phenyl is substituted by alkyl,  
alkoxy,



-63-


- halogen, or  
trifluoromethyl,  
heteroaryl,  
heteroaryl substituted by one to two substituents selected from the  
group consisting of:  
alkyl, or  
halogen,  
biphenyl,  
biphenyl substituted by alkyl,  
alkoxy,  
halogen,  
trifluoromethyl, or  
cyano,  
biphenylalkyl or  
biphenylalkyl wherein biphenyl is substituted by  
alkyl,  
alkoxy,  
halogen,  
trifluoromethyl, or  
cyano;  
D is zero or an integer of 1 to 3;  
L is zero or an integer of 1 to 3;  
R<sup>1</sup> is hydrogen  
a side chain or a natural amino acid or  
a side chain of an unnatural amino acid;  
Y is OR<sup>3</sup> wherein R<sup>3</sup> is hydrogen,  
methyl,  
ethyl, or  
benzyl, or  
NH-OR<sup>4</sup> wherein R<sup>4</sup> is hydrogen,  
alkyl, or  
benzyl;



-64-

and corresponding isomers thereof; or a pharmaceutically acceptable salt thereof.

2. The compound according to Claim 1 which is



- 5 3. The compound according to Claim 2 wherein  
n is zero or an integer of 1; and

X is -O-, or

-CH<sub>2</sub>-.

- 10 4. The compound according to Claim 3 wherein  
n is zero or an integer of 1;

X is -O-, or

-CH<sub>2</sub>-; and

R is hydrogen.

- 15 5. The compound according to Claim 4 wherein  
n is zero or an integer of 1;

X is -O-, or

-CH<sub>2</sub>-;

R is hydrogen; and

Y is OH.

- 20 6. The compound according to Claim 5 wherein  
n is zero or an integer of 1;

X is -O-, or

-CH<sub>2</sub>-;

R is hydrogen; and

- 25 Y is NHOH.



7. A compound selected from the group consisting of:

(S) 3-Methyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;

5 (S) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(S) 4-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;

(S) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-succinic acid;

10 (S) Phenyl-[(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)]-acetic acid;

(S) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-3-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-propionic acid;

15 (S) 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionic acid;

(S) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-4-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-butyric acid;

(S) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-5-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-pentanoic acid;

20 (S) 5-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid;

(S) 4-Phenylmethanesulfinyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;

25 (S) 4-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;

(S) 5-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid;

(S) 6-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-hexanoic acid;

30 (S) 7-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-heptanoic acid;

(S) 8-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-octanoic acid;



-66-

- (S) 4-Phenylsulfamoyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- (S) 4-Phenylmethanesulfonyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- 5 (S) 4-Benzylsulfanyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- (S) 3-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionic acid;
- (S) 4-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- 10 (S) 5-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid;
- (S) 6-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-hexanoic acid;
- 15 (S) 7-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-heptanoic acid;
- (S) 8-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-octanoic acid;
- (S) 2-(2,3-Dihydro-1H-8-oxa-cyclopenta[a]indene-6-sulfonylamino)-3-methyl-butyrac acid;
- 20 (S) 3-Methyl-2-(6,7,8,9-tetrahydro-5H-10-oxa-benzo[a]azulene-2-sulfonylamino)-butyric acid;
- (S) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;
- 25 (S) N-Hydroxy-3-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-succinamic acid;
- (S) N-Hydroxy-2-phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-acetamide;
- (S) N-Hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-3-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-propionamide;
- 30 (S) 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-N-hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionamide;



-67-

(S) N-Hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-4-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-butyramide;

5 (S) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-5-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-pentanoic acid hydroxyamide;

(S) 5-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid hydroxyamide;

10 (S) N-Hydroxy-4-phenylmethanesulfinyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(S) 4-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-N-hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(S) 5-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid hydroxyamide;

15 (S) 6-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-hexanoic acid hydroxyamide;

(S) 7-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-heptanoic acid hydroxyamide;

20 (S) 8-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-octanoic acid hydroxyamide;

(S) 4-Benzylsulfonyl-N-hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(S) N-Hydroxy-4-phenylsulfonyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

25 (S) N-Hydroxy-4-phenylmethanesulfonyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(S) N-Hydroxy-3-(1H-indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionamide;

30 (S) N-Hydroxy-4-(1H-indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(S) 5-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid hydroxyamide;



-68-

(S) 6-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-hexanoic acid hydroxyamide;

(S) 7-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-heptanoic acid hydroxyamide;

5 (S) 8-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-octanoic acid hydroxyamide;

(S) 2-(2,3-Dihydro-1H-8-oxa-cyclopenta[a]indene-6-sulfonylamino)-N-hydroxy-3-methyl-butyramide;

10 (S) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-5H-10-oxa-benzo[a]azulene-2-sulfonylamino)-butyramide;

(S) 3-Methyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyric acid;

(S) 3-Methyl-2-(9-methyl-6,7,8,9-tetrahydro-5H-carbazole-2-sulfonylamino)-butyric acid;

15 (S) 4-Phenyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyric acid;

(S) 4-Phenyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyric acid;

20 (S) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyramide;

(S) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyramide;

(S) N-Hydroxy-3-methyl-2-(9-methyl-6,7,8,9-tetrahydro-5H-carbazole-2-sulfonylamino)-butyramide;

25 (S) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyramide;

(S) 3-Methyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyric acid;

30 (S) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyramide;

(R) 3-Methyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;



-69-

(R) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(R) 4-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;

5 (R) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-succinic acid;

(R) Phenyl-[(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)]-acetic acid;

10 (R) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-3-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-propionic acid;

(R) 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionic acid;

(R) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-4-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-butyric acid;

15 (R) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-5-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-pentanoic acid;

(R) 5-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid;

20 (R) 4-Phenylmethanesulfinyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;

(R) 4-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;

(R) 5-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid;

25 (R) 6-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-hexanoic acid;

(R) 7-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-heptanoic acid;

30 (R) 8-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-octanoic acid;

(R) 4-Phenylsulfamoyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;



-70-

- (R) 4-Phenylmethanesulfonyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- (R) 4-Benzylsulfonyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- 5 (R) 3-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionic acid;
- (R) 4-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyric acid;
- (R) 5-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid;
- 10 (R) 6-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-hexanoic acid;
- (R) 7-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-heptanoic acid;
- 15 (R) 8-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-octanoic acid;
- (R) 2-(2,3-Dihydro-1H-8-oxa-cyclopenta[a]indene-6-sulfonylamino)-3-methyl-butyric acid;
- (R) 3-Methyl-2-(6,7,8,9-tetrahydro-5H-10-oxa-benzo[a]azulene-2-sulfonylamino)-butyric acid;
- 20 (R) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;
- (R) N-Hydroxy-3-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-succinamic acid;
- 25 (R) N-Hydroxy-2-phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-acetamide;
- (R) N-Hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-3-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-propionamide;
- 30 (R) 3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-N-hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionamide;



-71-

(R) N-Hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-4-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-butyramide;

5 (R) 2-(6,7,8,9-Tetrahydro-dibenzofuran-3-sulfonylamino)-5-(3,4,4-trimethyl-2,5-dioxo-imidazolidin-1-yl)-pentanoic acid hydroxyamide;

(R) 5-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid hydroxyamide;

10 (R) N-Hydroxy-4-phenylmethanesulfinyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(R) 4-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-N-hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(R) 5-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid hydroxyamide;

15 (R) 6-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-hexanoic acid hydroxyamide;

(R) 7-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-heptanoic acid hydroxyamide;

20 (R) 8-Phenyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-octanoic acid hydroxyamide;

(R) 4-Benzylsulfonyl-N-hydroxy-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(R) N-Hydroxy-4-phenylsulfamoyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

25 (R) N-Hydroxy-4-phenylmethanesulfonyl-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(R) N-Hydroxy-3-(1H-indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-propionamide;

30 (R) N-Hydroxy-4-(1H-indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-butyramide;

(R) 5-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-pentanoic acid hydroxyamide;



-72-

(R) 6-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-hexanoic acid hydroxyamide;

(R) 7-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-heptanoic acid hydroxyamide;

5 (R) 8-(1H-Indol-3-yl)-2-(6,7,8,9-tetrahydro-dibenzofuran-3-sulfonylamino)-octanoic acid hydroxyamide;

(R) 2-(2,3-Dihydro-1H-8-oxa-cyclopenta[a]indene-6-sulfonylamino)-N-hydroxy-3-methyl-butyramide;

10 (R) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-5H-10-oxa-benzo[a]azulene-2-sulfonylamino)-butyramide;

(R) 3-Methyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyric acid;

(R) 3-Methyl-2-(9-methyl-6,7,8,9-tetrahydro-5H-carbazole-2-sulfonylamino)-butyric acid;

15 (R) 4-Phenyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyric acid;

(R) 4-Phenyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyric acid;

20 (R) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyramide;

(R) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyramide;

(R) N-Hydroxy-3-methyl-2-(9-methyl-6,7,8,9-tetrahydro-5H-carbazole-2-sulfonylamino)-butyramide;

25 (R) N-Hydroxy-4-phenyl-2-(6,7,8,9-tetrahydro-dibenzothiophene-3-sulfonylamino)-butyramide;

(R) 3-Methyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyric acid; and

30 (R) N-Hydroxy-3-methyl-2-(6,7,8,9-tetrahydro-5H-fluorene-2-sulfonylamino)-butyramide;

and corresponding isomers thereof; or a pharmaceutically acceptable salt thereof.



-73-

8. The method of inhibiting a matrix metalloproteinase comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.
- 5 9. The method of inhibiting gelatinase A comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.
- 10 10. The method of inhibiting stromelysin-1 comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.
- 10 11. The method of inhibiting collagenase-3 comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.
- 15 12. The method of preventing atherosclerotic plaque rupture comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.
13. The method of inhibiting aortic aneurysm comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.
- 20 14. The method of inhibiting heart failure comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.
15. The method of preventing restenosis comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.



-74-

16. The method of controlling periodontal disease comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.
- 5 17. The method of treating corneal ulceration comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.
18. The method of treating burns comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.
- 10 19. The method of treating decubital ulcers comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.
20. The method of treatment for healing wounds comprising administering to a host suffering therefrom a therapeutically effective amount of a compound  
15 according to Claim 1 in unit dosage form.
21. The method of treating cancer comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.
22. The method of treating arthritis comprising administering to a host  
20 suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.
23. The method of treating osteoporosis comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.
- 25 24. The method of treating autoimmune or inflammatory disorders dependent upon tissue invasion by leukocytes comprising administering to a host



suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.

5           25.    The method of treating multiple sclerosis comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.

26.    The method of treating inflammation and pain comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.

10       27.    The method of treating acute and chronic neurodegenerative disorders selected from the group consisting of: stroke, head trauma, spinal cord injury, Alzheimer's disease, amyotrophic lateral sclerosis, cerebral amyloid angiopathy, AIDS, Parkinson's disease, Huntington's diseases, prion diseases, myasthenia gravis, and Duchenne's muscular dystrophy comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.

15

28.    The method of treating renal disease comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.

20       29.    The method of treating left ventricular dilation comprising administering to a host suffering therefrom a therapeutically effective amount of a compound according to Claim 1 in unit dosage form.


30.    The pharmaceutical composition comprising a compound according to Claim 1 in admixture with a pharmaceutically acceptable excipient, diluent, or carrier.



-76-

31. The pharmaceutical composition comprising a therapeutically effective amount of a compound according to Claim 1 in admixture with a pharmaceutically acceptable excipient, diluent, or carrier.
32. A method for preparing a compound of Formula Ie

5



Ie

wherein n is zero or an integer of 1 or 2;

X is -O-,

-S(O)<sub>p</sub> wherein p is zero or an integer of 1 or 2,

-N- wherein R<sup>2</sup> is hydrogen,

10

alkyl,  
|  
R<sup>2</sup> acyl, or  
benzyl,

-CH<sub>2</sub>, or

-C-; and

15




R<sup>1</sup> is hydrogen,

a side chain of a natural amino acid or a side chain of an unnatural amino acid; and corresponding isomers thereof; or

20

a pharmaceutically acceptable salt thereof which comprises treating a compound of formula (15)



(15)



-77-

wherein Ph is phenyl and n, X, and R<sup>1</sup> are as defined above with a base in a solvent to give a compound of Formula Ie and if desired, converting a compound of compound of Formula Ie to a corresponding pharmaceutically acceptable salt by conventional means and, if so desired, converting the corresponding pharmaceutically acceptable salt to a compound of Formula Ie by conventional means.



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/12273

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D307/91 C07D333/76 C07D307/93 A61K31/34 A61K31/38  
C07D405/04 C07D209/88 C07C311/21

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 09957 A (WARNER LAMBERT CO ;PICARD JOSEPH ARMAND (US); SLISKOVIC DRAGO ROBE) 12 March 1998 (1998-03-12) abstract; claims page 27 -page 33; examples	1,8-31
Y	WO 98 09934 A (BRIEN PATRICK MICHAEL O ;WARNER LAMBERT CO (US); PICARD JOSEPH ARM) 12 March 1998 (1998-03-12) abstract; claims page 33 -page 34 page 35 -page 36; examples	1,8-31
Y	WO 98 16503 A (AMERICAN CYANAMID CO) 23 April 1998 (1998-04-23) abstract; claims 1,6-11 page 8, line 14 - line 23	1,8-31
-/-		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"A" document member of the same patent family

Date of the actual completion of the international search

13 October 1999

Date of mailing of the international search report

22/10/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Paisdor, B



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/12273

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 20 33 959 A (GEIGY, J. R., A.-G.) 4 February 1971 (1971-02-04) page 2 page 34 -page 38; claims 10-15; examples 4-7	1
A	EP 0 555 824 A (DR. KARL THOMAE GMBH, GERMANY) 18 August 1993 (1993-08-18) abstract; claims 1,4,7	1,8



# INTERNATIONAL SEARCH REPORT

I. International application No.

PCT/US 99/12273

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 8-29  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 8-29  
are directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/12273

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9809957 A	12-03-1998	AU 4177097 A EP 0929542 A HR 970474 A	26-03-1998 21-07-1999 31-12-1998
WO 9809934 A	12-03-1998	AU 4159597 A CA 2256716 A EP 0931045 A	26-03-1998 12-03-1998 28-07-1999
WO 9816503 A	23-04-1998	AU 5145898 A EP 0938471 A	11-05-1998 01-09-1999
DE 2033959 A	04-02-1971	AT 299177 B BE 753172 A CH 517064 A CH 517688 A ES 381577 A ES 381578 A ES 381579 A ES 381580 A ES 381581 A ES 381582 A ES 381583 A FR 2059498 A NL 7009808 A US 3743663 A ZA 7004720 A	15-05-1972 08-01-1971 31-12-1971 15-01-1972 16-11-1972 16-11-1972 16-11-1972 16-11-1972 16-01-1973 16-01-1973 16-11-1972 04-06-1971 12-01-1971 03-07-1973 31-03-1971
EP 0555824 A	18-08-1993	DE 4204270 A AU 663556 B AU 3296893 A CA 2089466 A FI 930620 A IL 104703 A JP 6016648 A MX 9300735 A NZ 245886 A PL 297716 A US 5391556 A ZA 9300975 A	04-11-1993 12-10-1995 19-08-1993 14-08-1993 14-08-1993 13-07-1997 25-01-1994 01-08-1993 21-12-1995 24-01-1994 21-02-1995 12-08-1994



